



Traumatic And Knowledge on Self Efficacy During The Postpartum Recovery Period

Bina Melvia Girsang^{*1} , Eqlima Elfira¹ , Evi Indriani Br. Karo¹, Ismayadi¹

¹ Faculty of Nursing, Universitas Sumatera Utara, Medan, Indonesia

*Corresponding Author: binamelvia@usu.ac.id

ARTICLE INFO

Article history:

Received 31st May 2023

Revised 17th July 2023

Accepted 28th July 2023

Available online

<https://talenta.usu.ac.id/IJNS>

E-ISSN: 2685-7162

How to cite: Girsang, B.M. Eqlima, E., Evi, I. Br Karo. (2023). Traumatic And Knowledge On Self Efficiency During The Postpartum Recovery Period. *Caring: Indonesian Journal of Nursing Science*, 5(1), 34-40.

ABSTRACT

Information provision and postpartum care that is less than optimal is a problem that is often overlooked. The number of respondents based on the calculation of the sample size was 150 people, while the number of samples in this study were 134 postpartum mothers. The sample selection was carried out according to the probability sampling technique. Postpartum mothers were surveyed using a questionnaire about the history of childbirth trauma, the knowledge of postpartum mothers about self-care during the postpartum period, and the level of self-efficacy of postpartum mothers. The results of the study showed several factors that could affect the mother's self-efficacy in postpartum recovery, namely age (26.14 (4.96); $p = 0.00$), traumatic experience during childbirth (1.86 (0.34); $p = 0.000$), and mother's occupation (1.597 (0.49); $p = 0.02$), and knowledge (4.38 (1.68); $p = 0.000$). Nurses need to provide prenatal health education to increase women's beliefs about their behavior during labor and postpartum recovery.

Keyword: Education; Postpartum; Self-efficacy



This work is licensed under a
Creative Commons Attribution-
ShareAlike 4.0 International.

<https://doi.org/10.32734/ijns.v5i1.11404>

1. Introduction

Postpartum mothers are very vulnerable and can suffer from various complications, including mental illness that can affect their mental health achievement and reduce self-confidence after giving birth (Bahadoran et al., 2007). Maternal trauma during childbirth is most often associated with bad experiences such as psychological stress and trauma to the baby (Creedy et al., 2000). Maternal self-control tends to reduce pain during labor. In addition, good self-efficacy was associated with increased maternal satisfaction during the process (Christiaens & Bracke, 2007). The successful achievement of the mother's role refers to the mother's self-efficacy from the perspective of the ability to exert effects and mother's self-awareness in caring for the baby (Rahayuningsih, 2015).

Self-confidence can increase self-efficacy, but low self-confidence is often correlated with maternal fear of childbirth, causing anxiety. In addition, anxiety is highly correlated with efficacy and less effective (Schwartz et al., 2015). Trauma during childbirth has a significant impact on the mother's mental health and therefore affects her self-efficacy (Brazeau et al., 2018). Positive birth experiences affect postpartum comfort, mother psychology, and mother-infant relationships (Halperin et al., nd). Labor trauma is defined as an event that occurs during labor and the puerperium and results in serious or real or threatening injury or death to the mother or baby. Mother experiences serious fear, loss of control, discomfort, or vulnerability. Some

mothers feel that they are losing their self-esteem (Klein et al., 2014; Practitioner & 2014, nd; Research & 2015, 2015; Türkmen et al., 2020). One of the problems of postpartum stress is the lack of attention to the mother's emotional state (Razurel, C., Bruchon-Schweitzer, M., Dupanloup, A., Irion, O., & Epiney, 2011). The provision of information and postnatal care that is less than optimal is a problem that is often ignored because mothers are involved in complaints to health workers. Another thing that makes mothers uncomfortable is reporting their problems to health workers (Kline, CR, Martin, DP, & Deyo, 1998). In general, self-efficacy theory is a part of self-awareness that has a significant impact on a person's daily life. Knowledge of postnatal care enables postpartum mothers to successfully pass the important postnatal phases (Mirzaee, K., & Taghi Shakeri, 2015).

2. Method

The subjects of the study were 142 postpartum mothers in the working area of the Medan Sungal Health Center and the Medan Amplas working area. The number of respondents based on the calculation of the sample size was 150 people, while the number of samples in this study were 134 postpartum mothers. In this study, the criteria for postpartum mothers who were used as respondents were those with second-degree episiotomy wounds without labor complications. The exclusion criteria applied included the degree of episiotomy wound above 2 accompanied by complications of delivery. The sample selection was carried out according to the probability sampling technique. The questionnaire in this study was developed by researchers to assess postpartum mothers were surveyed using a questionnaire about the history of childbirth trauma, the knowledge of postpartum mothers about self-care during the postpartum period, and the level of self-efficacy of postpartum mothers. Filling out the questionnaire was done online using the Google form. Evaluation and revision model trials were carried out with a pre-post test using a T-test (Mother's occupation, episiotomy, and traumatic experiences during childbirth). In this study, the Pearson correlation test was also carried out on several components of the research variables such as occupation, level of education, and baby's weight.

3. Result and Discussion

The consistency of Bandura's self-efficacy theory, the most important predictor of successful behavior change, is self-efficacy in the ability to perform a behavior and the expectation that certain goals are achieved, including the principle of self-efficacy. Social cognitive theory states that exercise management, pain, fatigue, relaxation techniques, depression, diet, and communication with medical professionals are problem solving at the level of one's self-efficacy ability in setting goals (Munce, SE, Fehlings, MG, Straus, SE, Nugaeva, N., Jang, E., Webster, F., & Jaglal, 2014).

The ability of postpartum mothers to achieve self-efficacy during the postpartum recovery period is influenced by how knowledge the mother has, and not based on her level of education as stated in research conducted by Inu Martina (2018), which suggests that education level predictions have an effect on postpartum mother's self-efficacy. (Martina, 2018) and the results of other studies state that labor satisfaction is not related to a person's level of education (Ramie, A., Afiyanti, Y., & Pujasari, 2014).

Table. 1. Research Variable Analysis

Variable	Frequency	%	M (SD)	95% CI (Min-Max)	Statistical Test	p-Value
Age	134	100%	26.14 (4.96)	(25.29±26.98)	t-test	0.00*
Work						
Housewife	54	40.3	1,597 (0.49)	(1.51±1.68)	χ^2	0.025*
working mom	80	59.7				
Education Level						
Primary school	8	6	2,410 (0.62)	(2.30 ± 2.51)	Pearson correlation	0.1
Junior high school	65	48.5				
Senior High School	59	44.0				
College	2	1.5				
Baby BB						

Variable	Frequency	%	M (SD)	95% CI (Min-Max)	Statistical Test	p-Value
2500-3000	31	23.1	2,059 (0.72)	(1936±1832)	Pearson correlation	0.88
3001-3500	64	47.8				
3501-4000	39	29.1				
Episiotomy Procedure						
Done	93	69.4	1.306 (0.46)	(1.22 ± 1.38)	χ^2	0.24
Are not done	41	30.6				
Traumatic feeling during childbirth						
Traumatic	18	13.4	1.86 (0.34)	(1,807±1,924)	χ^2	0.00*
Not Traumatic	116	86.6				

In this study, several factors that can affect the mother's self-efficacy in postpartum recovery are age (26.14 (4.96); $p = 0.00$), traumatic experience in childbirth (1.86 (0.34); $p = 0.000$), and mother's occupation (1.597 (0.49); $p = 0.02$) where working mothers allow mothers to get more information from the work environment and the ability to access the information they need. Studies also show that self-efficacy will be achieved with good results if the mother has a good history of childbirth experience (Sánchez-Cunqueiro et al., 2018). Maturity of pregnant women is closely related to their willingness to give birth. Most pregnant women are in the age group that is not at risk of pregnancy, namely pregnant women aged 20-35 years are assumed to be relatively ready to face childbirth. Maternal care is expected to help reduce maternal anxiety when facing childbirth (Kartini, 2019). An explanation of the analysis of these factors can be seen in table 1.

The research statement states that the effect of knowledge on creating a sense of self-efficacy has not been thoroughly investigated (Schwartz et al., 2015). The results of previous studies stated that training is needed in dealing with childbirth and preparing for the postpartum period since pregnancy, this is believed to increase self-confidence, feelings, and a sense of empowerment in the mother (Byrne et al., 2014; Serçekuş et al., 2010). The results of other studies also prove that the knowledge possessed by the mother will help the mother in preparing for birth and good postpartum recovery because the mother's self-efficacy is influenced by the mother's perception of her delivery (Ip et al., 2009).

Table 2. The Effect of Knowledge on Postpartum Mother's Self-Efficacy

M (SD)	95% CI (Min-Max)	Statistical Test	p-Value
4.38 (1.68)	(26.71±27.46)	t-test	0.00*

Table 2 proves that knowledge affects the level of mother's self-efficacy in the postpartum recovery period (p -Value < 0,05). The information obtained by the mother has an impact on increasing delivery satisfaction, self-efficacy related to childbirth, control, and mother's level of knowledge (Bahrami et al., nd; Byrne et al., 2014; Miquelutti et al., 2013). In fact, while previous ineffectiveness of prenatal education about childbirth experiences has been reported (Fabian et al., 2005) there was no difference found among those taught prenatal courses. (Aba et al., nd). However, in several studies there is controversy about knowledge that can have an impact on knowledge and anxiety, self-control, labor pain, social and emotional adaptation of maternal self-efficacy, anxiety, stress, depression and childbirth is not clear and the evidence is inadequate (Aba et al., nd; Gagnon et al., nd).

Self-management and empowerment are important parts of the basic concepts in the treatment of self-efficacy (Duran et al., nd; Serçekuş et al., nd). Pregnant women with high self-efficacy during childbirth were found to be happy with their delivery (Berentson-Shaw et al., 2009). Self-efficacy is also related to the level of fear of childbirth. Poor self-efficacy in pregnant women reduces self-confidence during childbirth and has a negative impact on the active role of pregnant women during labor (Beebe, KR, Lee, KA, Carrieri Kohlman, V., & Humphreys, 2007; Salomonsson et al., 2013). Many studies report that low self-efficacy in pregnant women increases the perception of pain during childbirth, increases levels of anxiety and stress and develops into post-traumatic stress disorder during the puerperium (Salomonsson et al., 2013; Toohill et al., 2014).

4. Conclusion

The knowledge factor plays an important role in preparing mothers not only to face the postpartum recovery period, but also helps mothers to be confident in facing childbirth, and have positive coping to interpret a prosperous delivery. Knowledge affects the level of mother's self-efficacy in the postpartum recovery period. The information provided can include information about the physiology of birth exercise, breathing exercises, relaxation exercises, partner and professional support, self-efficacy, both during pregnancy and childbirth helps increase the perception ability to control, and facilitate the realization of effective coping strategies. Nursing professionals need to provide prenatal education to increase women's beliefs about their behavior during labor and postpartum recovery.

Acknowledgements

This research is fully supported by the research institute of the University of North Sumatra. The researcher is grateful for the opportunity given to carry out this research until it is completed and entered the publication stage.

Conflict of Interest

There is no conflict of interest in writing and submitting the publication of this research article.

References

- Aba, Y., research, NK-A. nursing, & 2017, undefined. (nd). Antenatal education on pregnant adolescents in Turkey: prenatal adaptation, postpartum adaptation, and newborn perceptions. Elsevier. Retrieved January 2, 2022, from <https://www.sciencedirect.com/science/article/pii/S1976131717301603>
- Bahadoran, P., Abbasi, F., Yousefi, AR, & Kargarfard, M. (2007). Evaluating the Effect of Exercise on the Postpartum Quality of Life. *Iranian Journal of Nursing and Midwifery Research* Winter, 12(1). www.mui.ac.ir
- Bahrami, N., Simbar, M., & SB-I. journal of fertility, & 2013, undefined. (nd). The effect of prenatal education on mother's quality of life during the first year postpartum among iranian women: a randomized controlled trial. *Ncbi.Nlm.Nih.Gov.* Retrieved January 2, 2022, from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3914493/>
- Beebe, KR, Lee, KA, Carrieri-Kohlman, V., & Humphreys, J. (2007). The effects of childbirth self efficacy and anxiety during pregnancy on prehospitalization labor. *Journal of Obstetrics, Gynecologic & Neonatal Nursing*, 36(5), 410–418.
- Berentson-Shaw, J., Scott, KM, & Jose, PE (2009). Do self-efficacy beliefs predict the primiparous labor and birth experience? A longitudinal study. *Journal of Reproductive and Infant Psychology*, 27(4), 357–373. <https://doi.org/10.1080/02646830903190888>
- Brazeau, N., Reisz, S., Jacobvitz, D., & George, C. (2018). UNDERSTANDING THE CONNECTION BETWEEN ATTACHMENT TRAUMA AND MATERNAL SELF-EFFICACY IN DEPRESSED MOTHERS. *Infant Mental Health Journal*, 39(1), 30–43. <https://doi.org/10.1002/IMHJ.21692>
- Byrne, J., Hauck, Y., Fisher, C., Bayes, S., & Schutze, R. (2014). Effectiveness of a mindfulness-based childbirth education pilot study on maternal self-efficacy and fear of childbirth. *Journal of Midwifery and Women's Health*, 59(2), 192–197. <https://doi.org/10.1111/JMWH.12075>
- Christiaens, W., & Bracke, P. (2007). Assessment of social psychological determinants of satisfaction with childbirth in a cross-national perspective. *BMC Pregnancy and Childbirth*, 7. <https://doi.org/10.1186/1471-2393-7-26>
- Creedy, DK, Shochet, IM, & Horsfall, J. (2000). Childbirth and the development of acute trauma symptoms:

- Incidence and contributing factors. *Birth*, 27(2), 104–111. <https://doi.org/10.1046/J.1523-536X.2000.00104.X>
- Duran, E., Dergisi, . A.-GT, & 2011, undefined. (nd). Kadınların sezaryen/vajinal doğuma ilişkin bakış açılarının analytical calibative. Researchgate.Net. <https://doi.org/10.13140/RG.2.1.2427.5287>
- Fabian, HM, Rådestad, IJ, & Waldenström, U. (2005). Childbirth and parenthood education classes in Sweden. Women's opinion and possible outcomes. *Acta Obstetricia et Gynecologica Scandinavica*, 84(5), 436–443. <https://doi.org/10.1080/J.0001-6349.2005.00732.X>
- Gagnon, A., Systematic, JS-C. database of, & 2007, undefined. (nd). Individual or group antenatal education for childbirth or parenthood, or both. *Cochranlibrary.Com*. Retrieved January 2, 2022, from <https://www.cochranlibrary.com/cdsr/doi/10.1002/14651858.CD002869.pub2/abstract>
- Halperin, O., Sarid, O., Midwifery, JC-, & 2015, undefined. (nd). The influence of childbirth experiences on women's postpartum traumatic stress symptoms: A comparison between Israeli Jewish and Arab women. Elsevier. Retrieved December 27, 2021, from <https://www.sciencedirect.com/science/article/pii/S0266613815000728>
- Ip, WY, Tang, CSK, & Goggins, WB (2009). An educational intervention to improve women's ability to cope with childbirth. *Journal of Clinical Nursing*, 18(15), 2125–2135. <https://doi.org/10.1111/J.1365-2702.2008.02720.X>
- Kartini, F. (2019). Developing Indonesian version of childbirth self-efficacy inventory (Cbsei)-C32. *Bali Medical Journal (Bali Med J)*, 8(3), 679–686.
- Klein, M., Vanderbilt, D., & Kendall-Tackett, K. (2014). PTSD and Breastfeeding: Let It Flow. *Infant, Child, and Adolescent Nutrition*, 6(4), 211–215. <https://doi.org/10.1177/1941406414541665>
- Kline, CR, Martin, DP, & Deyo, RA (1998). Health consequences of pregnancy and childbirth as perceived by women and clinicians. *Obstetrics & Gynecology*, 92(5), 842–848.
- Martina, I. (2018). THE EFFECT OF MATERNAL SELF EFFICACY ON THE PSYCHOLOGICAL WELL-BEING OF MATERNAL POST-DELIVERY SECTIO CAESARIA. <https://eprints.umm.ac.id/44765/>
- Miquelutti, MA, Cecatti, JG, & Makuch, MY (2013). Evaluation of a birth preparation program on lumbopelvic pain, urinary incontinence, anxiety and exercise: A randomized controlled trial. *BMC Pregnancy and Childbirth*, 13. <https://doi.org/10.1186/1471-2393-13-154>
- Mirzaee, K., & Taghi Shakeri, M. (2015). Maternal knowledge on postpartum care in healthcare centers of Mashhad, Iran in 2013. *Journal of Midwifery and Reproductive Health*, 3(4), 456–464.
- Munce, SE, Fehlings, MG, Straus, SE, Nugaeva, N., Jang, E., Webster, F., & Jaglal, SB (2014). Views of people with traumatic spinal cord injury about the components of self-management programs and program delivery: a Canadian pilot study. *BMC Neurology*, 14(1), 1–11.
- Practitioner, CZ-TN, & 2014, undefined. (nd). PTSD after childbirth: early detection and treatment. *Journals.Lww.Com*. Retrieved December 27, 2021, from https://journals.lww.com/tnpj/FullText/2014/03000/PTSD_after_childbirth__Early_detection_and.9.aspx
- Ramie, A., Afyanti, Y., & Pujasari, H. (2014). Self-Control and Self-Efficacy Increase Maternal Satisfaction Undergoing the Labor Process. *J Ners*, 9(1), 97–103.
- Razurel, C., Bruchon-Schweitzer, M., Dupanloup, A., Irion, O., & Epiney, M. (2011). Stressful events, social support and coping strategies of primiparous women during the postpartum period: a qualitative study. *Midwifery*, 27(2), 237–242.

- Research, CB-GQN, & 2015, undefined. (2015). Middle range theory of traumatic childbirth: The ever-widening ripple effect. *Journals.Sagepub.Com*, 2015, 1–13. <https://doi.org/10.1177/2333393615575313>
- Salomonsson, B., Gullberg, M., ... SA-... of P., & 2013, undefined. (2013). Self-efficacy beliefs and fear of childbirth in nulliparous women. *Taylor & Francis*, 3(34), 116–121. <https://doi.org/10.3109/0167482X.2013.824418>
- Sánchez-Cunqueiro, M., ... MC-JNE, & 2018, undefined. (2018). On the relation of self-efficacy and coping with the experience of childbirth. *Researchgate.Net*, 8(6). <https://doi.org/10.5430/jnep.v8n6p48>
- Schwartz, L., Toohill, J., Creedy, DK, Baird, K., Gamble, J., & Fenwick, J. (2015). Factors associated with childbirth self-efficacy in Australian childbearing women. *BMC Pregnancy and Childbirth*, 15(1). <https://doi.org/10.1186/S12884-015-0465-8>
- Serçekuş, P., Midwifery, HO-, & 2009, undefined. (nd). Fears associated with childbirth among nulliparous women in Turkey. *Elsevier*. Retrieved January 2, 2022, from <https://www.sciencedirect.com/science/article/pii/S0266613807000423>
- Serçekuş, P., Review, SM-IN, & 2010, undefined. (2010). Turkish women's perceptions of antenatal education. *Wiley Online Library*. <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1466-7657.2009.00799.x>
- Toohill, J., Fenwick, J., Gamble, J., Creedy, DK, Buist, A., Turkstra, E., & Ryding, EL (2014). A Randomized Controlled Trial of a Psycho-Education Intervention by Midwives in Reducing Childbirth Fear in Pregnant Women. *Birth*, 41(4), 384–394. <https://doi.org/10.1111/BIRT.12136>
- Türkmen, H., YALNIZ DİLCEN, H., & zçoban, FA (2020). Traumatic childbirth perception during pregnancy and the postpartum period and its postnatal mental health outcomes: a prospective longitudinal study. *Journal of Reproductive and Infant Psychology*, 1–13. <https://doi.org/10.1080/02646838.2020.1792429>