



Determination of Air Stability Parameter Threshold Value for Cumulonimbus and Thunderstorm Cloud Events at Kualanamu Meteorological Station

Rajab Prima*, Fendy Arifianto, Yosafat Donni H, and Avrionesti

State College of Meteorology Climatology and Geophysics, Tangerang Selatan, 15221, Banten, Indonesia.

*Corresponding Author: rajab.prima@stmkg.ac.id

ARTICLE INFO

Article history:

Received 22 June 2023

Revised 18 August 2023

Accepted 25 August 2023

Available online 31 August 2023

E-ISSN: 2656-0755

P-ISSN: 2656-0747

How to cite:

R. Prima, F. Arifianto, Y. Donni H, and Avrionesti, "Determination of Air Stability Parameter Threshold Value for Cumulonimbus and Thunderstorm Cloud Events at Kualanamu Meteorological Station," Journal of Technomaterial Physics, vol. 05, no. 02, pp. 99-103, Aug. 2023, doi: 10.32734/jotpv5i1.12487.

ABSTRACT

Many studies have carried out calculations related to atmospheric lability as a reference in weather forecasts, especially cumulonimbus clouds, and thunderstorms. However, many air lability index values are found to be inappropriate in each region because conditions in each region are different from each other in the region. So it is necessary to use precise index thresholds to determine weather conditions. In the study, observational data and data from Showalter Index (SI), Lifted Index (LI), K Index (KI), Severe Weather Threat Index (SWEAT), and Convective data were used. Available Potential Energy (CAPE) for ten years (2013-2022), then statistical calculations and verification for one year (2022) are carried out. The results obtained are the atmospheric stability index with the best accuracy in predicting the presence of cumulonimbus clouds and thunderstorms at the Kualanamu Meteorological Station, Deli Serdang is the best LI index to predict TS 00 and TS 12, and the best KI index to predict CB 00 and CB 12.

Keywords: Cumulonimbus, Thunderstorm, Stability index, Threshold, Meteorological Station

ABSTRAK

Banyak penelitian telah melakukan perhitungan terkait labilitas atmosfer sebagai acuan dalam prakiraan cuaca, khususnya awan cumulonimbus, dan badai petir. Namun, banyak ditemukan nilai indeks labilitas udara yang tidak sesuai di setiap wilayah karena kondisi di setiap wilayah berbeda satu sama lain di wilayah tersebut. Sehingga perlu menggunakan ambang batas indeks yang tepat untuk menentukan kondisi cuaca. Dalam penelitian tersebut digunakan data observasi dan data dari Showalter Index (SI), Lifted Index (LI), K Index (KI), Severe Weather Threat Index (SWEAT), dan data Konvektif. Energi Potensial Tersedia (CAPE) selama sepuluh tahun (2013-2022), kemudian dilakukan perhitungan statistik dan verifikasi selama satu tahun (2022). Hasil yang diperoleh adalah indeks kestabilan atmosfer dengan akurasi terbaik dalam memprediksi keberadaan awan cumulonimbus dan badai petir di Stasiun Meteorologi Kualanamu, Deli Serdang adalah indeks LI terbaik untuk memprediksi TS 00 dan TS 12, serta indeks KI terbaik untuk memprediksi CB. 00 dan CB 12.

Kata Kunci: Cumulonimbus, Badai Petir, Indeks Stabilitas, Threshold, Stasiun Meteorologi



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International.
<http://doi.org/10.32734/jotpv5i1.12487>

1. Introduction

Based on the recording of the Lightning Detection device at the Class 1 Geophysical Station Deli Serdang, there were 20,390 thunderstorm phenomena in the last week or the period from October 26 to November 1, 2020, caused by unstable atmospheric conditions due to strong convection and convergence, resulting in the lifting of air masses which can then expand vertically to create convective clouds, One of them is cumulonimbus clouds that produce thunderstorms [1].

Many air lability index values are not appropriate in a region because the conditions of each region are different from one region [1], so it is necessary to calculate a threshold that has high accuracy in predicting the presence of cumulonimbus and thunderstorm clouds well in the region for an index that characterizes the potential formation of cumulonimbus and thunderstorm clouds (SI, LI, KI, SWEAT, and CAPE) [1].

Based on this background, this study was conducted to find the threshold values of the SI, LI, KI, SWEAT, and CAPE indices to find out the best index that can be used to predict the presence of cumulonimbus clouds and thunderstorms in the Deli Serdang area.

2. Method

2.1. Data

The data processing in this study is upper air and surface air observation data at the time of thunderstorms and cumulonimbus clouds from the Kualanamumu Meteorological Station for ten years from 2013 to 2022. Top air observation data were collected during thunderstorms and cumulonimbus clouds at 00 UTC and observations at 12 UTC to see SI, LI, KI, SWEAT, and CAPE index values during thunderstorms and cumulonimbus clouds.

The synoptic data used for the analysis of thunderstorm events and cumulonimbus clouds is the current weather code 7www1w2 with passwords 13, 17, 91, 92, 93, 94, 95, 96, 97, 98, and 99, while for cumulonimbus cloud analysis using the 8NhClCmCh section CL password with passwords 3 and 9 [2].

2.2. Processing

The data processing techniques in this study are as follows:

- 1) The upper erial observation data was processed using the RAOB 5.7 application to obtain SI, LI, KI, SWEAT, and CAPE index values [3].
- 2) Data from observations of thunderstorms and cumulonimbus clouds were then grouped with RAOB 5.7 results according to time. Then separated by periods December, January, February (DJF), March, April, May (MAM), June, July, August (JJA), and September, October, and November (SON) [4-7].
- 3) Determine the threshold value according to the Sturges Equation which given in equation (1) to (3) [8,9]:

$$k = 1 + 3.322 \log \log n \quad (1)$$

$$R = X_n - X_t \quad (2)$$

$$i = \frac{(R)}{(k)} \quad (3)$$

where k is number of class intervals, n is amount of observational data, r is range, X_n is big data, X_t is small data, and i is class interval length.

- 4) Verify the value of the air lability parameter index for weather phenomena using the new Threshold. Verification is carried out using 2022 data. In verification, a 2×2 contingency table is used to see the level of accuracy of weather predictions.

Tabel 1. Contingency forecast table - results of observations [8].

	OBSERVATIONS			TOTAL
	Yes	No	TOTAL	
FORECAST	Yes	Hits (H)	False Alarms (FA)	Forecast 'Yes'
	No	Misses (M)	Correct Negatives (CN)	Forecast 'No'
	TOTAL	Forecast 'Yes'	Forecast 'No'	TOTAL

3. Results and Discussion

3.1. Threshold Value

Table 2. Threshold value of atmospheric stability index of Kualanamu Meteorological Station December, January, February (DJF).

	Index				
	LI	SWEAT	KI	SI	CAPE
TS 00	-1.98	190.01	34.6	-0.12	0
TS 12	-3.13	209.9	34.4	-0.08	942.33
CB 00	-2.38	191.71	32.9	-1.32	0
CB 12	-3.21	205.7	35.1	-0.78	316.4

Table 3. Threshold value of atmospheric stability index of Kualanamu Meteorological Station March, April, May (MAM).

	Index				
	LI	SWEAT	KI	SI	CAPE
TS 00	-3.15	200.6	32.6	-1.61	628.6
TS 12	-3.46	200.71	34.3	-1.54	699.4
CB 00	-2.75	202.81	34.1	-1.11	540.4
CB 12	-3.76	207.81	34.7	-0.88	1223.4

Table 4. Threshold value of atmospheric stability index of Kualanamu Meteorological Station June, July, August (JJA).

	Index				
	LI	SWEAT	KI	SI	CAPE
TS 00	-4.05	200.1	35.9	-0.56	0
TS 12	-3.42	221.58	37.3	-2.27	822.8
CB 00	-3.77	205.1	34.7	-1.17	0
CB 12	-4.05	219.78	35.9	-2.06	0

Table 5. Threshold value of atmospheric stability index of Kualanamu Meteorological Station September, October, November (SON).

	Index				
	LI	SWEAT	KI	SI	CAPE
TS 00	-2.36	195	35	-0.13	0
TS 12	-3.67	198.7	34.1	-0.78	932.4
CB 00	-2.36	198.3	33.8	-0.57	0
CB 12	-3.07	180.02	33	-0.08	339

The threshold value when cumulonimbus clouds begin to occur at LI values < -4.05 to -2.36 ; according to [10,11], this value allows very strong storms, even accompanied by tornadoes. The threshold value when cumulonimbus clouds begin to appear at SWEAT values > 180.02 to 219.78 , this value is based on the category of Wirjohamidjojo and Swarinoto (2013) in the category of allowing cumulus cloud growth. Cumulonimbus clouds occur at KI threshold values > 32.9 to 35.9 in the SON period. The SI threshold value at the time of cumulonimbus clouds is $SI < -2.06$ to -0.08 ; this value has a weak level of volatility and the possibility of small storms [10]. The CAPE threshold value shows cumulonimbus clouds occur at $CAPE > 0$ to 1223.4 , where according to Zakir, CAPE values < 1000 indicate weak atmospheric instability, while CAPE values of $1000 - 2500$ indicate moderate atmospheric instability [10,12].

The threshold value of thunderstorm events began to occur at $LI < -3.67$ in the SON period up to -2.96 , where according [13–15], the value of $0 < LI < -2$ indicates that storms can occur and require good storm support mechanisms and $-3 < LI < -5$ indicates an unstable atmosphere and allows storms to occur. The threshold value for thunderstorm events began to occur in $SWEAT > 190.01$ to 221.58 ; this value is based on the categories of [12] and included in the category of enabling cumulus cloud growth. Thunderstorm events occur at KI threshold values > 32.6 to 34.3 in the MAM period, where according to Wirjohamidjojo and Swarinoto (2013), KI values in the range of $31-35$ allow thunderstorms with a percentage of events of 61 to 80% and KI with a range of $36-40$ allow thunderstorms to occur. Thunderstorms percentage is $81 - 90\%$. The SI threshold value at the time of a storm is $SI < -0.13$ to -2.27 ; this value has a weak level of volatility, according

to Zakir [10]. The CAPE threshold value indicates that storms occur at $CAPE > 0$ to 942.33, whereas according to Zakir, this value indicates weak atmospheric instability [9,10].

3.2. Verification




Figure 1. The accuracy value of the atmospheric stability index when cumulonimbus clouds are present at the Kualanamu Meteorological Station.

The verification results show the accuracy value of each atmospheric stability index. At 00 UTC during the cumulonimbus cloud, the CAPE index showed the highest accuracy value compared to other atmospheric stability indices with an accuracy value of 0.63, and the thunderstorms also had the same highest accuracy value, the CAPE index of 0.57. At 12 UTC for cumulonimbus cloud events, the highest accuracy index value was the KI index with an accuracy value of 0.67, and the thunderstorm event was the CAPE index with an accuracy value of 0.65.

4. Conclusion

The difference in threshold values during TS 00, TS12, CB 00, and CB 12 events in the period December, January, February (DJF), March, April, May (MAM), June, July, August (JJA), and September, October, November (SON) at the Meteorological Station Kualanamu Meteorological Station, Deli Serdang illustrates different index values influenced by atmospheric conditions in each period and atmospheric stability index with the best accuracy value in predicting the presence of cumulonimbus clouds and thunderstorms in Kualanamu Deli Serdang Meteorological Station is the best LI index for predicting TS 00 and TS 12. The KI index is best for predicting CB 00 and CB 12.

References

- [1] N. F. Zahroh, N. W. S. P. Dewi, and D. Harsanti, "Indeks Labilitas Udara untuk Memprediksi Kejadian Badai Guntur pada Periode Puncak Musim Hujan Tahun 2016," *Jurnal Sains & Teknologi Modifikasi Cuaca*, vol. 18, no. 1, 2017, doi: 10.29122/jstmc.v18i1.1764.
- [2] M. Budiarti, M. Muslim, and Y. Ilhamsyah, "Studi Indeks Stabilitas Udara Terhadap Prediksi Kejadian Badai Guntur (Thunderstorm) di Wilayah Stasiun Meteorologi Cengkareng Banten," *Jurnal Meteorologi dan Geofisika*, vol. 13, no. 2, 2012, doi: 10.31172/jmg.v13i2.125.
- [3] B. Primohadi Syahputra, M. Yusuf, Y. Donni Haryanto, and N. Florida, "Visualization of Aerological Diagram and Analysis of Atmospheric Sounding Information Using Raob with Model Data During Low Visibility Conditions at Cengkareng Meteorological Station," *Proceedings of International Conference on Multidisciplinary Research*, vol. 5, no. 2, 2022, doi: 10.32672/pic-mr.v5i2.5420.
- [4] A. A. Rusdin, P. Palloan, S. Subaer, and A. Prasetyo, "Uji Akurasi Ambang Batas Indeks Stabilitas Atmosfer Terhadap Pembentukan Thunderstorm dan Awan Cumulonimbus di Stasiun Meteorologi Kelas I Sultan Hasanuddin," *Jurnal Fisika Unand*, vol. 12, no. 2, 2023, doi: 10.25077/jfu.12.2.268-274.2023.
- [5] S. Sukarna, W. Sanusi, and S. D. R. Ningsih, "Model Vector Autoregressive Exogenous dan Aplikasinya pada Curah Hujan Kota Makassar," *Journal of Mathematics Computations and Statistics*, vol. 2, no. 2, 2020, doi: 10.35580/jmathcos.v2i2.13745.

- [6] B. HK. Tjasyono, *Meteorologi Indonesia Volume I : Karakteristik dan Sirkulasi Atmosfer*, vol. I. 2012.
- [7] E. Aldrian, "Meteorologi Laut Indonesia," *Researchgate*, no. June, 2008.
- [8] I. J. A. Saragih, B. W. Yonas, and N. Rinaldy, "Uji Ambang Batas Baru Indeks Total-Total Untuk Prakiraan Kejadian Awan Cumulonimbus Dan Thunderstorm Di Bandara Kualanamu," in *Seminar Nasional Fisika dan Aplikasinya*, Jatinangor: Universitas Padjadjaran, 2017, pp. 203–210.
- [9] D. S. Wilks, *Statistical methods in the atmospheric sciences, second edition*, vol. 14, no. 2. 2007.
- [10] A. Zakir, W. Sulistya, and M. K. Khotimah, *Perspektif Operasional Cuaca Tropis*. 2010.
- [11] Y. Ilhamsyah, "Kajian pendahuluan analisis peramalan thunderstorm untuk penyusunan indeks dasar adaptasi kegiatan pertambakan (Suatu tinjauan meteorologi di Jakarta)," *Depik*, vol. 1, no. 1, 2013, doi: 10.13170/depik.1.1.27.
- [12] A. Arsali, O. C. Satya, S. Supardi, and I. Purna, "Penentuan Koefisien untuk Perhitungan Suhu Udara Rata-Rata Harian Data Stasiun Klimatologi Palembang," *Jurnal Meteorologi dan Geofisika*, vol. 16, no. 1, 2015, doi: 10.31172/jmg.v16i1.260.
- [13] L. O. Bangsawan and B. Tjasyono HK, "Kajian Ambang Batas Indeks Stabilitas Udara Terkait Kejadian Hujan Lebat Dan Badai Guntur Di Makasar," *STMKG*, 2015.
- [14] K. Isnoor and H. A. Rejeki, "Analisis Kondisi Atmosfer Saat Terjadi Borneo Vortex dan Pengaruhnya Terhadap Curah Hujan di Wilayah Kalimantan Barat dan Kepulauan Riau (Periode 2006 – 2016)," *Seminar Nasional Penginderaan Jauh ke-5*, 2018.
- [15] J. A. I. Paski, A. Sepriando, and D. A. S. Pertiwi, "Pemanfaatan Teknik RGB pada Citra Satelit Himawari-8 untuk Analisis Dinamika Atmosfer Kejadian Banjir Lampung 20 - 21 Februari 2017," *Jurnal Meteorologi Klimatologi dan Geofisika*, vol. 4, no. 3, 2019, doi: 10.36754/jmkg.v4i3.48.