

# Relationship between the Practise of Consuming Pork Lawar and the Prevalence of Taeniasis in Kerobokan Village, Badung Regency, Bali

Ni Luh Putu Ayu Gayatri<sup>1\*</sup>, Hemma Yulfi<sup>2</sup>

<sup>1</sup>Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia

<sup>2</sup>Department of Parasitology, Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia

**Abstract. Introduction:** *Taeniasis is an infection brought on by Taenia sp. Tapeworms, one of which is Taenia solium, whose intermediate host is swine. Adult tapeworms develop in the intestines of the host (human) after consuming infected pork that was not properly prepared. Lawar babi is one of the Balinese dishes that utilises cooked, undercooked, or even uncooked pork. Balinese, including residents of Kerobokan Village, are accustomed to eating pork lawar. Objective:* To determine the relationship between the consumption of pork lawar and the incidence of taeniasis in the village of Kerobokan, Badung Regency, Bali. **Methods:** This is a correlative categorical research with a cross sectional design. The study population is people in Kerobokan Village, Bali Province. Using power of 0.8 strength of linear correlation of 0.25, and additional 10% for incomplete data, we obtain sample size of 128 people who meet inclusion criteria. Data was obtained by filling out a set of validated questionnaire. **Results:** Sixty-five out of 128 respondents (50.8%) were found to possess a positive inclination towards the consumption of pork lawar, and only 1 (one) respondent (0.08%) had a history of taeniasis. The significance value of the correlation ( $p$ ) between the habit of eating pork lawar and a history of taeniasis was 0.312. **Conclusion:** There is a lack of correlation between the practise of consuming pork lawar and the prevalence of taeniasis among the population residing in Kerobokan Village, located in the Badung Regency of Bali.

**Keywords:** Pork lawar, Taenia solium, Taenia saginata, Taeniasis

Received: 11 August 2023 | Revised: 31 August 2023 | Accepted: 31 August 2023

## 1 Introduction

Taeniasis is caused by *Taenia solium* when raw or undercooked pork is consumed [1]. Tapeworm infection is associated with numerous clinical symptoms. The eggs of tapeworms excreted in the faeces of human carriers are infectious to swine. *Taenia solium* eggs can also infect humans if

they are ingested via the faecal-oral route or by consuming contaminated food or water, resulting in tissue infection with parasitic larvae [2].

*Taenia* infection in adult worms is diagnosed by the presence of gravid segments in a stool sample. Tissue cysts in worm larvae can be identified via biopsies, radiography (calcified cysts), and computed tomography (brain cysts). The sensitivity and specificity of serology utilising the indirect hemagglutination test and ELISA are beneficial but variable. The 2010 Global Burden of Disease (GBD) Survey estimated that human taeniasis and cysticercosis caused by *Taenia solium* result in 503,000 (95% confidence interval [CI]: 379,000–663,000) disability-adjusted life years (DALYs) lost annually. Extrapolating DALYs associated with epilepsy from the 2010 GDB survey reveals that the DALYs associated with *Taenia solium* are approximately 2.7 million (95% CI: 2.16-3.6 million). *Taenia solium* is also believed to be responsible for up to 28,000 fatalities annually (95% CI: 21,000–36,000) [3][4].

Indonesia is a tropical region with a significant population infected with taeniasis, and the prevalence of cysticercosis and taeniasis ranges between 2% and 48% [5]. North Sumatra, Bali, and Papua are the three most prevalent endemic regions. In Indonesia, the spread of tapeworms is largely influenced by the cultural and religious context of each island. One of them is the island of Bali, which is home to many popular local and tourist dishes, such as pork lawar. This cuisine contains pork that is cooked, undercooked, or raw [6]. Consuming pork lawar fresh or undercooked can increase the risk of intestinal tapeworm infection. Therefore, this study aimed to see the relationship between intestinal taeniasis and the habit of pork lawar consumption in Bali.

## 2 Method

This is a cross-sectional study, focused on correlational analysis within categorical variables. The study was carried out during the period of July to November 2022. The study population consisted of the people residing in Kerobokan Village, located in the Badung Regency of Bali Province. A total of 128 patients were enrolled in the study using the sequential sampling method, with a statistical power of 0.8, a linear correlation strength of 0.25, and an additional 10% inclusion to account for potential missing data. The process of data collecting involved the completion of online questionnaires that had undergone rigorous testing to ensure their validity and reliability. The data was subjected to analysis using the SPSS statistical package for Windows version 25.0, developed by IBM Corp. in Armonk, NY. The study included both univariate and bivariate techniques, specifically employing Spearman's rho correlation coefficient.

## 3 Result

The 128 questionnaires collected were examined for completeness. The data was subsequently presented in tabular format, providing information on the distribution of sociodemographic

characteristics, the distribution of risk factors, the distribution of taeniasis cases, and a cross-tabulation of both variables.

**Table 1** Sociodemographic distribution of respondents

Characteristics	Frequency (n)	Percentage (%)
<b>Gender</b>		
Male	55	43
Female	73	57
<b>Age (years)</b>		
<30	59	46,1
30-50	55	43
51-70	11	8,6
>70	3	2,3
<b>Last education</b>		
College	53	41,4
Senior high school	57	44,5
Junior high school	9	7
Elementary school	6	4,7
No school	3	2,3
<b>Type of work</b>		
Government employees	8	6,3
Private employees	25	19,5
Self-employed	45	35,2
Farmer	2	1,6
Housewife	17	13,3
Student	18	14,1
No work	12	9,4
<b>Total</b>	<b>128</b>	<b>100</b>

**Table 2** Distribution of risk factors

Characteristics	Frequency (n)	Percentage (%)
<b>Ownership of livestock</b>		
Pigs	13	10,2
Cattle	8	6,3
Others	40	31,3
None	67	52,3
<b>Maintenance of pig farm</b>		
Good	12	92,3
Moderate	1	7,7
Bad	0	0
<b>Habit of consuming pork lawar</b>		
Good	65	50,8
Bad	63	49,2
<b>Habit of processing pig meet</b>		
Good	5	3,9
Moderate	123	96,1
Bad	0	0
<b>Total</b>	<b>128</b>	<b>100</b>

Taeniasis is made based on the respondent's history of expelling proglottids from the anal voluntarily or involuntarily (thru defecation).

**Table 3** Distribution of taeniasis

Characteristics	Frequency (n)	Percentage (%)
Negative history of expelling proglottids	127	99,2
Positive a history of expelling proglottids	1	0,8
<b>Total</b>	<b>128</b>	<b>100</b>

The bivariate analysis conducted using Spearman's rho test yielded a significance value of 0.312 ( $p > 0.05$ ), indicating that there is no statistically significant link between the consumption of pork lawar and the occurrence of taeniasis.

**Table 4** The relationship between the habit of consuming pork lawar and the incidence of taeniasis

Variable		Incident taeniasis						Score <i>P</i>
		Negative history		Positive history		Total		
		n	%	N	%	N	%	
The habit of consuming pork lawar	Good	65	100	0	0	65	100	0,312
	Bad	62	98,4	1	1,6	63	100	
Total		127	99,2	1	0,8	128	100	

#### 4 Discussion

The majority of participants indicated that they did not possess agricultural land. A small proportion of individuals possessed pigs or animals. The majority of pig farmers demonstrated a high level of farm management, as seen by their diligent maintenance of their farms. This included the frequent cleaning of pig pens and the provision of specialised animal feed to ensure the well-being of the pigs. This finding suggests that the participants demonstrated a strong knowledge of effective agricultural practises, particularly in the domain of pig farming [7]. A study conducted in Denpasar, Bali, revealed that pig farms in the area have been effectively managed and maintained. These farms have adhered to proper protocols for pig rearing, ensuring the animals' protection from parasite illnesses [8]. This phenomenon may also be attributed to the extensive health promotion efforts conducted by healthcare professionals, village leaders, and other extension organisations or community groups.

Based on the observed dietary behaviour pertaining to the consumption of pork lawar, it was found that a majority of the participants continued to consume this dish in its raw or undercooked form. The cultural and customary practises of the Balinese people have an impact on the consumption of raw or undercooked pork lawar. Lawar is a customary culinary offering

originating from the Balinese culture, typically presented during significant Hindu festivities in Bali. However, it is not uncommon for lawar to be served on ordinary occasions as well [6].

The survey revealed that a significant proportion of participants demonstrated proper handling and preparation techniques for pork, hence ensuring compliance with hygiene regulations. Nevertheless, the community continues to offer pork in its raw or undercooked state, so presenting

a potential health risk. The observed phenomenon can be attributed to the pervasive impact of Balinese culture and customs [6][9]. During significant cultural observances like Galungan and Kuningan, it is customary for the community to make arrangements for the procurement and preparation of high-quality pigs to be ceremonially slaughtered [7]. Pork lawar is traditionally employed as a culinary component within ceremonial contexts. A diverse range of pig lawar preparations are presented as offerings to the deities, encompassing both cooked and uncooked variations. Following its utilisation as an offering, the food will subsequently be distributed and consumed collectively by the community.

Notwithstanding the aforementioned risk factors, the present study revealed that a significant proportion of participants did not possess a previous medical record of taeniasis. There was a singular individual with a documented medical history of taeniasis, who happened to be in the advanced stages of life, nearing the age of 100. This phenomenon can be comprehended by taking into account additional risk factors apart from the intake of raw pork lawar that were prevalent in the past. For instance, the limited utilisation of sanitary latrines and inadequate livestock management practises were also significant contributors. Historically, it was common practise to allow pigs and other livestock to roam freely or remain uncaged, primarily due to the abundance of available open space that might serve as a supplementary food source for these animals. During that period, there was a significant dearth of public knowledge regarding health matters, specifically in relation to taeniasis. Currently, the residents of Bali possess a heightened level of knowledge on taeniasis and its associated risk factors. The declining prevalence of taeniasis in Bali over time is evident. A study conducted in multiple districts and cities in Bali revealed a fluctuating prevalence of taeniasis over a period of time. The prevalence in Gianyar regency experienced a significant decline of sixfold over a four-year period, specifically from 2002 to 2006. A similar pattern was also observed in the regencies of Denpasar, Badung, and Karangasem. This is due to advancements in personal hygiene practises, enhanced management techniques in pig farming, and the spread of metropolitan regions. Consequently, individuals who formerly engaged in unrestricted livestock rearing are now compelled to either cease farming activities altogether or confine their free-range operations within enclosures [10]. A study conducted in a distinct city in Indonesia, with by comparable socio-demographic traits, revealed a complete absence of taeniasis cases in recent years. This outcome can likely be attributed to the population's practise of abstaining from the intake of raw or undercooked meat. The thorough cooking of meat to the point of being well done has the capacity to fully deactivate the cysticercus, hence eliminating the potential risk of contracting a taeniasis infection [11].

The present investigation did not identify any significant association between the practice of consuming pork lawar and the occurrence of taeniasis. In contrast, empirical investigations conducted in Peru have established a correlation between the consumption of pork and the

prevalence of taeniasis [12]. This is due to the ingestion of pork contaminated with the *Taenia solium* parasite. The porcine subjects in the research were reared in an environment devoid of enclosures, while the local populace had limited availability of sanitary amenities, particularly in terms of hygienic latrines. The presence of freely wandering swine and inadequate sanitary facilities is likely to heighten the susceptibility to *Taenia solium* infection [13].

Although there is a common practise of consuming raw meat, it is important to note that pigs that are not infected with *Taenia* sp. do not pose a risk of transmitting taeniasis. Pigs that are reared under appropriate conditions, housed in enclosures situated at a considerable distance from human dwellings, and are restricted from unrestricted movement, in addition to being provided with adequate nutrition, will yield pigs of superior grade [14].

## 5 Conclusion

Majority of people in Kerobokan Village, Badung Regency, Bali still consumed undercooked or raw pig meat. However, it is worth noting that a significant proportion of the individuals surveyed did not report any previous instances of taeniasis. There is a lack of correlation between the practise of consuming pork lawar and the prevalence of taeniasis in Kerobokan Village, located in the Badung Regency of Bali.

## REFERENCES

- [1] M. Coral-Almeida, S. Gabriel, E. N. Abatih, N. Praet, W. Benitez and P. Dorny, "Taenia solium Human Cysticercosis: A Systematic Review of Sero-epidemiological Data from Endemic Zones around the World," *National Library of Medicine*, 2015.
- [2] WHO, "Taeniasis/cysticercosis Fact Sheet," 2022. [Online]. Available: <https://www.who.int/news-room/fact-sheets/detail/taeniasis-cysticercosis>. [Accessed 15 Maret 2022].
- [3] A. L. Okello and L. F. Thomas, "Human taeniasis: current insights into prevention and management strategies in endemic countries," *Dovepress*, vol. 10, p. 107-116, 2017.
- [4] Torgerson PR, Devleeschauwer B, Praet N, Speybroeck N, Willingham AL, Kasuga F, et al. World Health Organization Estimates of the Global and Regional Disease Burden of 11 Foodborne Parasitic Diseases, 2010: A Data Synthesis. *PLoS Med* 2015;12.

- 
- [5] Wandra T, Depary AA, Sutisna P, Margono SS, Suroso T, Okamoto M, et al. Taeniasis and cysticercosis in Bali and North Sumatra, Indonesia. *Parasitol Int* 2006;55:S155–60. <https://doi.org/10.1016/j.parint.2005.11.024>.
- [6] L. Gutama, A. Salim, N. Jaya, I. Suyasa, M. Devinta and I. Wahyuniari, "Red Worm Free Program for Ada Village Communities," *Serving Udayana Bulletin*, vol. 20, 2021.
- [7] Trisdayanti NPE, Sawitri AAS, Sujaya IN. Sanitation Hygiene and the Potential Presence of E. Coli Virulence Genes in Lawar in Kuta: Challenges of Tourism and Food Health in Bali. *Public Health and Preventive Medicine Archive* 2015;3:99–105.
- [8] H. S. Bekti, N. Habibah, L. P. Rinawat, N. P. C. D. P. Yasa, O. D. G. Rindi, N. K. A. K. Dewi, N. P. A. D. Savitri and A. Rakhmawati, "Microscopic Identification of Taenia solium in Pig Farming," *Health Journal*, vol. 12, 2021.
- [9] Susanty E. Taeniasis solium and Sisticercosis in human. *Jurnal Ilmu Kedokteran (Journal of Medical Science)* 2019;12:1–6.
- [10] T. Wandra, S. S. Margono, M. S. Gafar, J. M. Saragih, P. Sutisna, A. R. Sudewi, A. Depary, H. Yulfi, H. Darlan, M. Okamoto, M. O. Sato, Y. Sako, M. Nakao, K. Nakaya, P. S. Craig and A. Ito, "Current Situation of Taeniasis and Cysticercosis in Indonesia," *Tropical Medicine and Health*, vol. 35, p. 323-328, 2007.
- [11] L. G. Barus, "Relationship of Taenia solium Infection to Pork Consumption in Simalingkar B Area," *Medan : University of North Sumatra*, 2021.
- [12] Wilson M, Sarti E, Roberts J, Gutierrez IO, Schantz PM, Flisser A, et al. Prevalence and Risk Factors for Taenia Solium Taeniasis and Cysticercosis in Humans and Pigs in a Village in Morelos, Mexico. *Am J Trop Med Hyg* 1992;46:677–85.
- [13] N. S. Watts, M. Pajuelo, T. Clark, M.-C. I. Loader, M. R. Verastegui, C. Sterling, J. S. Friedland, H. H. Garcia and R. H. Gilman, "Taenia solium Infection in Peru: A Collaboration between Peace Corps Volunteers and Researchers in a Community Based Study," *Plos One Collection Personalized Medicine*, 2014.
- [14] C. V. Simatupang, "Description of the State of Taenia solium in the Pardomuan Nauli Community, Selayang Village, Finish District, Langkat Regency," *Republic of Indonesia Ministry of Health Medan Health Polytechnic*, 2019.