



## Research Article

# Soil Characteristics of Settlements in Sukadono Hamlet, Helvetia District, Deli Serdang Regency

Muhammad Azran Bin Azmi<sup>\*1</sup>, Yoan Carolina Panggabean<sup>2</sup>, Yunilda Andriyani<sup>2</sup>  
Bambang Prayugo<sup>3</sup>

<sup>1</sup>Undergraduate Program of Medicine, Faculty of Medicine, Universitas Sumatera Utara, Medan, 20155, Indonesia

<sup>2</sup>Department of Parasitology, Faculty of Medicine, Universitas Sumatera Utara, Medan, 20155, Indonesia

<sup>3</sup>Department of Surgery, Faculty of Medicine, Universitas Sumatera Utara, Medan, 20155, Indonesia

### Abstract

**Background:** Soil is where humans can do an abundance of activities. Soil can be contaminated by various factors. Soil is also a medium of development, storage and transmission of several types of worms, commonly referred to as Soil-Transmitted Helminths (STH). **Objective:** The purpose of this study was to find out the characteristics of residential land in the Sukadono Hamlet Helvetia District Deli Serdang. **Methods:** The sample in this study was 72 soils taken from the front, back left or right of the house. The soil is measured in temperature, humidity, pH value and is checked using a Digital Soil Analyzer. **Results:** The results showed that most of the soil characteristics in Sukadono Hamlet Helvetia District Deli Serdang were abnormal. The STH examination results also showed that (45.8%) of the 72 soil samples examined positively contained STH. Hookworms were found in (32.9%) soil samples, *Ascaris lumbricoides* species in (5.3%) soil samples, *Strongyloides stercoralis* species were found positive in (7.9%) soil samples and a mixture of STH is present in 2 (3.9%) soil samples. **Conclusion:** The resident soil in research areas had been contaminated by Soil Transmitted Helminths.

**Keywords:** contamination, soil characteristics, soil transmitted helminth

Received: 01 February 2023 | Revised: 23 August 2023 | Accepted: 01 September 2023

## 1. Introduction

Soil contamination are generally caused by a group of nematodes referred to as Soil Transmitted Helminth (STH) [1]. STH causes infections in humans due to contact with eggs or larvae the are developing in warm and moist soil temperatures that is in tropical and subtropical Countries, STH consists of *Ascaris lumbricoides*, *Trichuris trichiura*, and hookworms (*Americanus Necator* and *Ancylostoma duodenale*) [2, 3]. Roundworm diseases is a disease that is caused by Soil Transmitted Helminths and are often suffered by citizens in developing countries and are estimated about more than 60% of the population. STH infection can occur either alone or simultaneously with some other type of worm at once because of low enviroment sanitation quality [4]. Human will most often be exposed to the infective form of STH in an area with poor sanitation quality [5].

More than 1.5 billion people or 24% of the world's population are infected with worms that transmitted through soil in the whole world. The infection spreads largely tropical and subtropical areas, with the biggest amount of infection happens in Sub-Saharan Africa, America, China and East Asia. More than 267 million preschool-aged children and more than 568 million school-age children live in areas where these parasites are transmitted intensively and need treatment and intervention prevention [6].

In Jakarta there is other research reported, *Ascaris lumbricoides* worm infection is as big as (62.2%), and (48%) for *Trichuris trichiura* and (0.72%) for hookworms [7]. Based on research by Samad in the city of Medan, indicated the presence of eggs STH as much as 52.5% of the 80 soil samples studied whilst the

\* Corresponding author at: Undergraduate Program of Medicine, Faculty of Medicine, Universitas Sumatera Utara, Medan, 20155, Indonesia.

E-mail address: robloxmeh@yahoo.com

Prevalence of STH Infection in State I-VI elementary students 105296 in the District Percut Sei Mr. Deli Serdang Regency, Sumatera Utara in 2019 is as much as (29.9%) [7].

Soil Transmitted Helminths can cause deficiency in nutrition, anemia, disruption of growth and cognitive development especially on children and are usually referred to as neglected disease due to generally afflicting the poor people in the world and gets less attention compared to other diseases [8]. Sukadono Hamlet, Helvetia District, Deli Serdang that is used as research is a resident housing area. The environment around the area tends to not be cleaned and appears to have a lot of rubbish and livestock feces. The villagers also work as scavengers and pig/chicken ranchers.

Based on the background above, this study aims to find out the characteristics of the soil in the Sukadono Hamlet, Helvetia District, Deli Serdang. The soil characteristics studied are include aspects of temperature, humidity, pH value, presence of STH and types of STH in the residential land of the Sukadono Hamlet Helvetia District, Deli Serdang.

## 2. Methods

The research type used is descriptive-observational with the cross-sectional method and was held in November 2022. Research is done on the residential soil that is in Sukadono Hamlet, Helvetia District, Deli Regency Serdang, Sumatera Utara. The examination of temperature, humidity and pH values is carried out using the Digital Soil Analyzer tool. The tip of the Digital Soil Analyzer is stabbed into the ground with a depth of 3 cm. Then the button is pressed and released. The results of the temperature, humidity and pH value will be displayed on the Digital Soil Analyzer. The microscopic examination of the soil was done in the Laboratory of Parasitology at the Faculty of Medicine, Universitas Sumatera Utara, located on Dr. Mansyur Road No 5, USU Campus. The population of this study is residential land of the villagers in Sukadono Hamlet, Helvetia District, Regency Deli Serdang, Sumatera Utara Province and is obtained by probability sampling with simple random type sampling because of homogenous population. The sampling period in this study was for 1 month.

Soil sample examination is carried out by putting 50 gr soil in a plastic pot that is already cleaned and 50 gr of soil is moved based on the plastic pot to a glass beaker, then add 250 milliliters of aquades & stir until its mixed using a stirrer. Then insert the mixed soil into a 50- size milliliter centrifuge tube using filter paper. Then, centrifuge the tube using a speed of 2,000rpm for two minutes and then the supernatant liquid is removed using a pipette then dissolve 0.5 milliliters of soil deposits using 150 milliliters of saturated magnesium sulfate solution in beakers, and stirred until it is mixed. Then pour the solution into a 15 milliliters centrifuge tube, then centrifuge the tube using the speed of 2,500 rpm for five minutes.

In addition, the centrifuged tubes with the centrifuged contents will be placed on a tube rack & a solution of magnesium sulfate is dripped until full or the surface looks like convex but does not spill. The top of the centrifuge tube is closed using cover glass once every 24 hours. Lift the cover glass in an upright and straight position then put in on glass object. Finally, examine the preparations under a microscope using 100x magnification & 400x to identify if there is or the not any Soil Transmitted Helminths.

This method is done to raise eggs / larvae Soil Transmitted Helminths to the surface cover glass to see the presence of STH in the study sample. Ethical Approval obtained from the Medical Ethics Committee of the Faculty of Medicine at the Universitas Sumatera Utara with Number 159/UN5.2.1.1.2.6/SPB/2022.

## 3. Results

In this study soil samples began with examinations using the Digital Soil Analyzer tool. The findings begin with an examination of the temperature of the soil sample. The normal temperature value of the used soil is 24.9°C -26.6° C and the rest are categorized as abnormal soil temperatures.

**Table 1.** Soil temperature frequency distribution

Characteristics temperature	Frequency	Percentage
Normal	22	31
Abnormal	50	69.4
Total	72	100

Based on the study of the author, The soil in the Sukadono Hamlet settlement, different temperatures were found recorded. Table 1.2 will show the average value of the temperature of soil sample.

**Table 2.** Table of average temperature values

Temperature	Frequency	Total
33°C	10	330
32°C	17	544
31°C	8	248
30°C	6	180
29°C	2	58
27°C	1	27
26°C	4	104
25°C	9	225
24°C	9	216
23°C	5	115
21°C	1	21

Based on Table 2, found a flat value of the average soil temperature is 29 ° C. In this study, soil samples began with examinations using the Digital Soil Analyzer tool. The examination is continued by recording the humidity of the soil sample. Dry soil was found in Sukadono Hamlet. Table 1.3 shows the frequency distribution of soil moisture.

**Table 3.** Soil temperature frequency distribution

Characteristics humidity	Frequency	Percentage
Dry	62	86
Moist	10	14
Total	72	100

Based on the results of table 3, it was found that 62 soil samples (86%) were dry, and 10 soil samples (14%) were moist. In this context, the research uses the Digital Soil Analyzer tool. The pH value of the soil is recorded. Soil pH values are categorized as normal and abnormal. Normal categories include pH values of 6 to 8 whilst abnormal pH values includes values below 6 and above 8. The overall pH value of soil samples is normal. Table 4 shows the frequency distribution of soil pH values.

**Table 4.** Frequency distribution of soil pH values

Characteristics pH value	Frequency	Percentage
Normal	72	100
Abnormal	0	14
Total	72	100

Based on the analysis from Table 1.4, it was found that the pH values of all 72 soil samples (100% ) were in normal pH values of 6-8 and were not found in soils where the pH values were abnormal. In this study, the author has found that 2 different soil pH values, 6 and 7 were found in Sukadono Hamlet settlement. Table 5 showed the average value of soil pH. Therefore in Table 5, the average value of the average soil pH is 7.

**Table 5.** Data table average value of pH value

pH value	Frequency	Total
6	1	6
7	71	497
Total	72	503

Microscopic examination in the laboratory is used for the purpose of finding STH in the soil sample. In this study 33 soil samples were found STH and 39 were not found STH. Table 6 shows the frequency distribution of STH presence.

**Table 6.** Distribution presence of STH

Absence of STH	Frequency	Percentage
STH found	33	45.8
STH not found	39	54.2
Total	72	100

Based on analysis from Table 6, it is found that on 33 samples (45.8%) found parasites, 39 samples (54.2%) not found STH. There are many types of STH that can be found in soil. In this study, the type of STH found were *Ascaris lumbricoides*, Hookworm, *Strongyloides stercoralis*, and a mixture of *Ascaris lumbricoides* and Hookworm or *Strongyloides stercoralis* and Hookworm. The common STH found in the soil is Hookworm. Table 7 will show the presence of different types of STH in soil samples.

**Table 7.** Presence of different types of STH in soil sample

Types of STH	Frequency	Percentage
<i>Ascaris lumbricoides</i>	4	5.3
Hookworm	25	32.9
<i>Strongyloides stercoralis</i>	6	7.9
Mixture	3	3.9
Negatif	39	50
Total	77	100

Based on the analysis from Table 7, *Ascaris lumbricoides* were found in 4 samples (5.3%), Hookworm in 25 samples (32.9%), *Strongyloides stercoralis* in 6 samples (7.9%), A mixture of STH were 2 in (3.9%) samples, and the latter 39 samples (50%) were found with no STH contamination.

#### 4. Discussion

In this study the temperature of residential soil in the Sukadono Hamlet, 22 (31%) soil samples were found in normal values i.e. in between 24.9°C-26.6°C. The results of this study differ from the research conducted by Oyewole and Simon-Oke [9], they recorded the temperature of the soil during the dry season is 35.77 °C and the rainy season is 27.69°C with a total range between 21°C - 41°C. The temperature of the soil in the Sukadono Hamlet Helvetia District Deli Serdang Regency is great for the growth of Soil Transmitted Helminths. Saharman et al and Handayati et al [10, 11] reported that the optimum temperature for growth of larvae and hookworm eggs (Hookworm) is between 23°C- 25°C for *Ancylostoma duodenale* and 28°C-32°C for *Necator Americanus*.

Characteristics of the soil that was recorded were dry, sandy, or gull soil. A total of 62 (86%) of the soil samples studied were dry soil and 10 (14%) of the examined soil were moist soil. The dry and sandy soil is suitable for larval growth of Hookworm which is widely found in the research sites. There is also warm and moist soil that is a perfect environment for *Ascaris lumbricoides* [12]. Moisture and soil are measured according to the research done by Unaeni (13) i.e. by using the Digital Soil Analyzer tool. But the results of this study differ from the results of Unaeni [13] which are shown that moist soils (0.955) are the most affected variables in the presence of STH eggs. The results of the study of soil pH values in the Sukadono Hamlet in total were 72 (100%) soil samples that were in normal pH values of 6 to 8. This result is in accordance with the research done by Oyewole and Simon-Oke [9] i.e. recorded soil pH ranges from 6.10 to 9.00.

Based on microscopic examination, it was found that in the entirety of the 72 samples, 39 samples were free from parasites, and 33 samples were found parasites. The parasites that were successfully identified were Filariform and rhabditiform larvae Hookworm, *Ascaris lumbricoides* larvae, Filariform larvae and rhabditiform *Ascaris lumbricoides*, *Ascaris lumbricoides* egg, Hookworm egg. This is similar with the research done by Natasya [14], where she found as many as 3 (3%) *Ascaris lumbricoides* eggs, 12 (12%) Hookworm eggs and a mixture of both *Ascaris lumbricoides* as well as Hookworm in 2 (2%) of the soil samples. The most parasitic finding is the discovery of Hookworm larvae which has been found in 31 (32.9%) soil samples. The results of the study are in line with research conducted by Apriyan & Song in 2018 [3] that uses the method Brine flotation for their research. Research found that Soil Transmitted Helminths in 13 of the total 92 samples discovered about around 90% of the soil sample which is 12 were

discovered to be Hookworm larvae. Hookworm found in the whole study were in the larvae form because the samples were not done straight but coveted more first because wanting to wait until all the sample were collected [3]. According to Soedarto's research [15] Hookworm eggs which are released with the stool will grow into their rhabditiform in 2 days up to a week and will develop into their filariform stage.

In the 72 soil samples collected, 4 (5.3%) were found species of eggs and larva of *Ascaris lumbricoides*. These results differ from studies from Steinbaum et al (12), which took 2107 samples from several places adjacent to nearby houses, 13.0% of the total soil sample was *Ascaris lumbricoides*. *Ascaris lumbricoides* found in this study are *Ascaris lumbricoides* larvae, *Ascaris lumbricoides* Corticated Infective eggs and *Ascaris lumbricoides* Corticated Non-Infective eggs.

No species of *Trichuris trichiura* (0%) were found in the entire soil sample in this research. Maybe because of unsuitable soil conditions such as clay, the growth of *Trichuris trichiura* cannot be done. The optimum condition for the development *Trichuris trichiura* is 30°C with clay type soil [16]. The temperature at the sample site were very optimum for the growth of *Trichuris trichiura* that is 30° C- 35°C however the type of soil that were found there were sandy and thus does not match with the optimum growth condition for *Trichuris trichiura* that develops well in more claylike soil.

In this study, filariform and rhabditiform larvae of *Strongyloides stercoralis* were also found in 6 (7.9%) soil samples. These results are in line with the research of Dr. Invoke and friends [17] who took samples from 797 patients containing those involved with geophagy, namely soil consuming activities. The research shows that 1% - 20% of the accumulated soil contains *Strongyloides stercoralis*. Maybe because the research location is not as dense, prevalence *Strongyloides stercoralis* were not much.

## 5. Conclusion

Based on the results of data analysis in this study, it can be concluded that the soil temperature in the residential area of Sukadono Hamlet, Helvetia District, Deli Serdang Regency is not within the normal range. The humidity in the area is classified as dry, while the soil pH is within the normal limits. The presence of Soil Transmitted Helminths was detected in the soil of the residential area, with identified types including *Ascaris lumbricoides*, Hookworm, and *Strongyloides stercoralis*.

## 6. Data Availability Statement

The datasets generated and analyzed during the current study are not publicly available due to privacy and ethical considerations but are available from the corresponding author upon reasonable request.

## 7. Ethical Statement

This study was approved by the Medical Ethics Committee Universitas Sumatera Utara under approval number: 159/UN5.2.1.1.2.6/SPB/2022.

## 8. Author Contributions

All authors contributed to the design and implementation of the research, data analysis, and finalizing the manuscript.

## 9. Funding

No funding.

## 10. Conflict of Interest

Authors declares no conflict of interest.

## References

- [1] Noviasuti AR. Infeksi soil transmitted helminths [Internet]. 2015 [cited 2022 Apr 22]. Available from: <https://jke.kedokteran.unila.ac.id/index.php/majority/article/view/1483/1322>.
- [2] Murni PH, Lubis M, Fujiati II. Hubungan infeksi soil transmitted helminths Dengan Kemampuan Kognitif, status Nutrisi, Dan Prestasi belajar pada anak sekolah dasar di Desa Sikapas kabupaten mandailing natal. *Sari Pediatri*. 2018;19(5):279. <https://doi.org/10.14238/sp19.5.2018.279-83>.
- [3] Apriyan F, Song C. Prevalensi soil-transmitted helminths di tanah taman perumahan Kecamatan Grogol, Cengkareng, Dan Kalideres Jakarta Barat. *Tarumanagara Medical Journal*. 2018;1(1):74–9. <https://doi.org/10.24912/tmj.v1i1.2519>.
- [4] Silva N da, Farhan A, Malatuzzzufa NI. Identifikasi Soil Transmitted Helminth (STH) pada feses Petani di Desa Plandi Kabupaten Jombang. 2020. <http://repo.itskesicme.ac.id/id/eprint/4064>.

- [5] Wandra T, Darlan DM, Yulfi H, Purba IE, Sato MO, Budke CM, et al. Soil-transmitted helminth infections and taeniasis on Samosir Island, Indonesia. *Acta Tropica*. 2019;202:105250. <https://doi.org/10.1016/j.actatropica.2019.105250>.
- [6] World Health Organization. Soil-transmitted helminth infections [Internet]. 2022 [cited 2022 Mar 27]. Available from: <https://www.who.int/news-room/fact-sheets/detail/soil-transmitted-helminth-infections>.
- [7] Tapiheru MJ, Zain N. Prevalensi infeksi soil transmitted Helminth Pada murid Sekolah Dasar negeri 105296 Kecamatan Percut Sei Tuan, kabupaten deli Serdang, Sumatera utara. *JIMKI: Jurnal Ilmiah Mahasiswa Kedokteran Indonesia*. 2021;8(3):1–7. <https://doi.org/10.53366/jimki.v8i3.249>.
- [8] Indriyati L, Juhairiyah, Hairani B, Fakhrizal D. Gambaran Kasus stunting Pada 10 desa di Kabupaten Tanah Bumbu tahun 2018. *Jurnal Kebijakan Pembangunan*. 2020;15(1):77–90. <https://doi.org/10.47441/jkp.v15i1.57>.
- [9] Oyewole OE, Simon-Oke IA. Ecological risk factors of soil-transmitted helminths infections in Ifedore District, Southwest Nigeria. *Bull Natl Res Cent*. 2022;46:24. <https://doi.org/10.1186/s42269-022-00700-8>.
- [10] Saharman S, Mayulu N, Hamel R. Hubungan Personal Hygiene Dengan Kecacingan Pada Murid Sekolah Dasar Di Kabupaten Bolaang Mongondow Utara. *J-Kp*. 2013;1(1). Available from: <https://ejournal.unsrat.ac.id/v3/index.php/jkp/article/view/2231>.
- [11] Handayani D, Ramdja M, Nurdianthi IF. Hubungan infeksi soil transmitted helminths (STH) Dengan Prestasi belajar Pada Siswa sdn 169 di Kelurahan Gandus Kecamatan gandus Kota Palembang [Internet]. 2015 [cited 2022 May 18]. Available from: <https://ejournal.unsri.ac.id/index.php/mks/article/view/2750>.
- [12] Steinbaum L, Mboya J, Mahoney R, Njenga SM, Null C, Pickering AJ. Effect of a sanitation intervention on soil-transmitted helminth prevalence and concentration in household soil: A cluster-randomized controlled trial and risk factor analysis. *PLoS Negl Trop Dis*. 2019;13(2):e0007180. <https://doi.org/10.1371/journal.pntd.0007180>.
- [13] Unaeni U. Kontaminasi Telur soil transmitted helminths (STH) pada buah stroberi di perkebunan Dan Pasar Ciwidey, Bandung Selatan, Jawa Barat [Internet]. Jakarta: UIN Syarif Hidayatullah; 2019 [cited 2022 Jun 18]. Available from: <https://repository.uinjkt.ac.id/dspace/handle/123456789/50310>.
- [14] Tesalonika NC. Identifikasi soil transmitted helminths pada tanah Peternakan domba di Desa Manggis kecamatan serbajadi kabupaten Serdang Bedagai [Internet]. Medan: Universitas Sumatera Utara; 2020 [cited 2022 Sept 21]. Available from: <https://repository.usu.ac.id/handle/123456789/46663>.
- [15] Soedarto. Buku ajar parasitologi kedokteran. 2nd ed. Jakarta: Sagung Seto; 2016.
- [16] Sastrawan IG, Setiabudi J, Sanjiwani NP, Indriyani NK, Laksemi DA. Risk factors of soil transmitted helminth infection among primary school students. *Health Sci J Indones*. 2020;11(2):126–32. <https://doi.org/10.22435/hsji.v11i2.2885>.
- [17] Ivoke N, Ikpor N, Ivoke O, Ekeh F, Ezenwaji N, Odo G, et al. Geophagy as risk behaviour for gastrointestinal nematode infections among pregnant women attending antenatal clinics in a humid tropical zone of Nigeria. *Afr Health Sci*. 2017;17(1):24–31. <https://doi.org/10.4314/ahs.v17i1.5>.