



SUMEJ
Sumatera Medical Journal

Journal homepage: <https://talenta.usu.ac.id/smj>



Parasite Contamination of Water Source at Pangkalan Masyhur Sub-district, Medan Johor District

Sunna Vyatra Hutagalung^{*1}, Rahmah Khairuna Damanik², Pongruj Rattaprasert³

¹Department of Parasitology, Faculty of Medicine, Universitas Sumatera Utara, Medan 20155, Indonesia

²Undergraduate Program in Medicine Faculty of Medicine, Universitas Sumatera Utara, Medan 20155, Indonesia

³Department of Protozoology, Faculty of Tropical Medicine, Mahidol University, Bangkok 10400, Thailand

*Corresponding Author: sunna.vyatra@usu.ac.id

ARTICLE INFO

Article history:

Received 16 April 2023

Revised 19 July 2024

Accepted 29 August 2024

Available online 5 September 2024

E-ISSN: 2622-1357

P-ISSN: 2622-9234

How to cite:

S. V. Hutagalung, R. K. Damanik, P. Rattaprasert, "Parasite Contamination of Water Source at Pangkalan Masyhur Sub-district, Medan Johor District," *Sumatera Medical Journal*, vol. 07, no. 03, September 2024

ABSTRACT

According to World Health Organisation (WHO) estimates, at least 1,8 billion people consume contaminated water worldwide, hence most of the diarrheal disease in the world is attributable to unsafe water, and the lack of sanitation and hygiene. Water utilization by humans and animals can generate biological pollutants which may contaminate the water. These contaminants come from waste products that may contain bacteria, protozoa, and other parasites. This was a descriptive study with cross sectional design and simple random sampling to determine the contamination of water source by pathogens, particularly parasites at Region V Pangkalan Masyhur sub-district, Medan Johor district, Medan, North Sumatera in 2012. Samples were obtained from each house, depending on the main water source used, either from tap water, open wells, or the nearby river streams. Modified Caldwell's sedimentation technique was applied to examine the samples under light microscope. Out of the 103 water samples collected, most were from tap water (78 samples), followed by open wells (19 samples), and only six were from the nearby river streams. We found *Paramecium caudatum* from nine samples (8,7%) collected from the open wells and the nearby river streams, followed by hookworm larva in one tap water sample (1%). No protozoa such as *Entamoeba* spp., *Cryptosporidium* spp., and *Giardia* spp. found in any of the samples. Based on this data, the abundant presence of *P.caudatum* represented the condition of high organic debris in the open wells and the nearby river streams. Additionally, the presence of hookworm larva in the tap water indicated the distribution system was contaminated with the surrounding soil. Hence it is imperative to provide counseling about optimal environmental sanitation and clean water management, including proper maintenance for water source and its distribution system. Tap water distribution, including its network of pipe deliverance system, must be checked and maintained regularly.

Keyword: Contamination, Parasites, Water



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International.

<http://doi.org/10.32734/sumej.v7i3.11826>

1. Introduction

Clean water is a primary need of human. WHO estimated at least 1.8 billion people use water source contaminated with feces, either from human and/or animal excreta. Contaminated water is linked to transmission of diseases, including diarrhea and dysentery. Contaminated drinking-water is estimated to cause 502,000 diarrheal deaths each year [1]. Diarrhea is mostly linked to contaminated food and water, plus lack on sanitation and hygiene.

The contaminated water may contain biologic pollutants from feces, such as virus, bacteria, and other parasites. According to WHO, among the important parasites known to contaminate water source are *Entamoeba histolytica*, *Cryptosporidium parvum*, and *Giardia duodenalis* [2]. *E. histolytica* is a pathogenic amoeba commonly distributed worldwide, causing amebic colitis and dysentery [3]. This protozoa infects human through fecal-oral route, and commonly obtained from water contaminated with human feces [4]. *Cryptosporidium* spp. is a parasitic protozoa known to infect the intestines of humans and various animals, such as cattle, sheep, rodents, cats, dogs, birds, fish, and reptiles. Transmission occurred by fecal-oral route through ingestion of infectious oocyst. In human, the species *Cryptosporidium hominis* and *Cryptosporidium parvum* are known to induce acute watery diarrhea, vomiting, nausea and fever in immunocompromised hosts, which can be life-threatening [5]. Outbreaks have been reported in water playgrounds and childcare settings [6]. The flagellate *Giardia* spp. is associated with diarrhea and malabsorption in human. The manifested diarrhea usually involved the appearance of foul-smelling, greasy feces, with abdominal discomfort [7]. The main route for infection is through infective cyst in contaminated food and water [8]. Improper chlorination and poor maintenance of water treatment system will maintain the oocyst/cyst stage of these parasites in the water; hence transmission may occur [9]. Additionally, other pathogenic parasites including helminths' larvae can also pollute water through contact with contaminated soil, further suggesting the poor state of the water purity.

This study aimed to determine the contamination of water source by pathogens, particularly parasites, at Region V Pangkalan Masyhur sub-district, Medan Johor district, Medan, North Sumatera in 2012. The region covered 5,5-hectare area, at the height of 2,5-37,5 meters above sea level, with the topography tends to crook to the north. This area has nearby river stream (Deli River), local small scales farming, and some of its residence still used open wells as water source. This situation creates the inclination for water source contamination by the parasites in the neighborhood.

2. Method

This was a descriptive study with cross-sectional design and simple random sampling. Samples were collected and examined from March to August 2012. The daily main water source samples, either from tap water or open wells or the nearby river streams, were collected from each house. The collection acquired one sample per each house. Sample size calculation at 95% confidence level and precision level $\pm 5\%$ at predicted population size of 140 houses was estimated as 103 samples. Water samples were collected from each of the house with a sterile plastic container, then examined immediately at Parasitology Laboratory, Faculty of Medicine, Universitas Sumatera Utara. Modified Caldwell's sedimentation technique was applied to examine the samples according to the protocol [10, 11]. In brief, 20 milliliter of each sample was centrifuged for 5 minutes at 2500 rpm. The supernatant was discarded. The remaining sediment was added with lugol, homogenized, then being directly examined under light microscope (Olympus® CX21, magnification 10x to 1000x). The conduction of the study was approved for ethical clearance from the Health Research Ethical Committee of Faculty of Medicine, Universitas Sumatera, by No. 196/KOMET/FKUSU/2012.

3. Result and Discussion

The main water sources for daily activities found in the region were as follows; 78 from tap/piped water from the local municipal waterworks, 19 from open wells, and six from the nearby river streams. From identification, we did not find any presence of *Entamoeba* spp., *Cryptosporidium* spp., or *Giardia* spp. as previously hypothesized (Table 1). We found ciliate *Paramecium caudatum* in nine samples (8.7%); six from the open wells' sources and three from the nearby river streams (Figure 1). Interestingly, we also found one sample from the tap water containing hookworm larva (1%), as shown in Figure 2.

Table 1. Direct microscopy examination result of the identified microorganisms found in the water samples

Water source	Total number	Parasite identified				
		<i>Entamoeba</i> spp.	<i>Cryptosporidium</i> spp.	<i>Giardia</i> spp.	<i>Paramecium caudatum</i>	<i>Hookworm</i> larva
Open wells	19	-	-	-	6	-
Tap water	78	-	-	-	-	1
Nearby river stream	6	-	-	-	3	-
Total	103	-	-	-	9	1

Figure 1. Ciliate *Paramecium caudatum* found in one of the samples.

Figure 2. Hookworm larva found in one sample.

Entamoeba spp. infection in human is commonly transmitted in areas with poor sanitation which allows contamination of drinking water and food with feces, with estimated prevalence range from 1% to 40% of the population in Central and South America, Africa, and Asia. In this study, we did not find any *Entamoeba* spp. in the samples. This may be due to the method used, because microscopic examination sensitivity for *Entamoeba* spp. detection varied from 10 to 50%, depending on the examiner's experience [12, 13].

Cryptosporidium spp. is an enteric protozoa found in both humans and livestock. Waterborne transmission for *Cryptosporidium* spp. has been reported via both surface water and groundwater [9, 14]. In this study, we also did not find any presence of *Cryptosporidium* spp. oocyst, probably due to the same reason for the negative findings on *Entamoeba* spp. Additionally, we only performed simple direct examination, without further stainings such as Giemsa stain or acid-fast Ziehl-Neelsen to increase the sensitivity, due to budget limitation.

Giardia spp. is one of the most prevalent intestinal protozoan flagellate of the human. The transmission of *Giardia* spp. occurs either by direct contact with the feces of infected mammals or by contaminated food and water. Although there were local small farms in the area, we did not find any *Giardia* spp. in the collected samples nearby. This is probably due the similar reason mentioned for detection of *Entamoeba* spp. and *Cryptosporidium* spp. The collection of water sample was only done once for each, and probably the presence of those parasites was low below the detection sensitivity level of simple microscopic examination [15]. We did not employ any specific antigen detection methods such as direct fluorescent antibody tests, ELISA, or other immunochromatographic assays to diagnose these parasites in this study.

In this study, the most common finding we obtained was the presence of *Paramecium caudatum*. This ciliated protist was found in water source samples from open wells and nearby river streams. *Paramecium caudatum* is ubiquitous in fresh and brackish water, usually in the sludge of organic matters or the sewage area. *Paramecia* feed on microorganisms like bacteria, algae, and yeasts, and they play an important role in wastewater biological purification processes [16, 17]. The presence of *Paramecium caudatum* indicated that the water source was full with sludge of organic matters and possible presence of abundant coliform bacteria, which may not be that clean or safe for daily use.

Interestingly, we also found one sample from tap/piped water contaminated with hookworm larva. Hookworm is nematode parasite commonly transmitted through infested soil. It may cause chronic gastrointestinal tract infection, sucking the host's blood which leads to iron deficiency anemia. Zoonotic hookworm larva can also produce cutaneous larva migrans. Hookworm larva usually contaminates soil [18]. This finding indicates that the piped water distribution system was likely leaked and under poor maintenance, exposing contact to the surrounding contaminated soil. The proper and regular maintenance of the water distribution, including the distributing pipe maintenance, is essential to ensure the quality of the delivered safe water.

Although the time of the study was considered dated, but the significance of the message derived from this is still quite relevant. Recent massive sewage drainage excavation, an on-going project in the whole Medan city, is a changing environment which may pose threat to the safe water distribution system by introduction of contaminants to pollute the water. The risk for broken pipes during excavation, and the exposure of the water to the excavation's dumps, may expose risks to safe water delivery system maintenance for the city's households.

4. Conclusion

From this study, we conclude that none of the water source samples collected from Region V Pangkalan Masyhur sub-district, Medan Johor district, Medan, North Sumatera contained any of the *Entamoeba* spp., *Cryptosporidium* spp., or *Giardia* spp. protozoa. The dominating finding was the presence of *Paramecium caudatum*. We also found one tap water sample contaminated with hookworm larva.

In this study, each sample was only obtained once from one random site of the main water source from each home. Additionally, we only employed direct method of microscopic observation with lugol, without further stainings. The method is simple, affordable, can easily be done in the field, and producing quick result. However, further studies with multiple sample collection and employment of more advanced methods are suggested to increase sensitivity and to obtain more representable results.

Ethics Approval: Sumatera Medical Journal (SUMEJ) is a peer-reviewed electronic international journal. This statement below clarifies ethical behavior of all parties involved in the act of publishing an article in Sumatera Medical Journal (SUMEJ), including the authors, the chief editor, the Editorial Board, the peer-reviewer and the publisher (TALENTA Publisher Universitas Sumatera Utara). This statement is based on COPE's Best Practice Guidelines for Journal Editors.

Authors Contributions: All authors contributed to the design and implementation of the research, data analysis, and finalizing the manuscript.

Funding: No funding.

Disclosure: Authors declares no conflict of interest.

References

- [1] World Health Organization, "Water," WHO | Regional Office for Africa, 2015. [Online]. Available: <https://www.afro.who.int/health-topics/water> [Accessed: April 15, 2023].
- [2] World Health Organization, Guidelines for drinking-water quality, 4th ed. Geneva: World Health Organization, 2022. Available: <https://iris.who.int/bitstream/handle/10665/352532/9789240045064-eng.pdf?sequence=1>.
- [3] C. Marie, W.A. Petri Jr., "Amoebic dysentery," *BMJ Clin Evid*, vol. 0918, 2013. PMID: 23991750; PMCID: PMC3758071.
- [4] Centers for Disease Control and Prevention, "Amebiasis," CDC, 2019. [Online]. Available: <https://www.cdc.gov/dpdx/amebiasis/index.html> [Accessed July 18, 2024].
- [5] European Centre for Disease Prevention and Control, "Cryptosporidiosis," ECDC, 2012. [Online]. Available: <https://www.ecdc.europa.eu/en/cryptosporidiosis> [Accessed July 18, 2024].
- [6] R. Gharpure, *et al.*, "Cryptosporidiosis Outbreaks — United States, 2009–2017," *MMWR. Morbidity and Mortality Weekly Report*, vol. 68, no. 25, pp. 568–572, 2019. <https://doi.org/10.15585/mmwr.mm6825a3>.
- [7] H. Hooshyar, *et al.*, "Giardia lamblia infection: review of current diagnostic strategies," *Gastroenterol Hepatol Bed Bench*, vol. 12, no. 1, pp. 3–12. 2019.
- [8] Centers for Disease Control and Prevention, "CDC - DPDx – Giardiasis," CDC, 2019. [Online]. Available: <https://www.cdc.gov/dpdx/giardiasis/index.html> [Accessed July 18, 2024].
- [9] F.E. Adeyemo, *et al.*, "Efficiency of chlorine and UV in the inactivation of Cryptosporidium and Giardia in wastewater," *PLoS One*, vol. 14, no. 5, e0216040, 2019. <https://doi.org/10.1371/journal.pone.0216040>.
- [10] M. Camacho, *et al.*, "Lutz's spontaneous sedimentation technique and the paleoparasitological analysis of sambaqui (shell mound) sediments," *Memórias do Instituto Oswaldo Cruz*, vol. 108, no. 2, pp. 155–159, 2013. <https://doi.org/10.1590/0074-0276108022013005>.
- [11] F. C. Caldwell and E. L. Caldwell, "Preliminary Report on Observations on the Development of Ova of Pig and Human Ascaris under Natural Conditions, and Studies of Factors Influencing Development," *The Journal of parasitology*, vol. 14, no. 4, pp. 254–254, 1928. <https://doi.org/10.2307/3271383>.
- [12] H. Uslu, O. Aktas, M.H. Uyanik, "Comparison of Various Methods in the Diagnosis of Entamoeba histolytica in Stool and Serum Specimens," *Eurasian J Med*, vol. 48, no. 2, pp. 124–129, 2016. <https://doi.org/10.5152/eurasianjmed.2015.0074>.
- [13] M. Tanyuksel M., W.A. Petri Jr., "Laboratory diagnosis of amebiasis," *Clin Microbiol Rev*, vol. 16, no. 4, pp. 713–729, 2003. <https://doi.org/10.1128/CMR.16.4.713-729.2003>.
- [14] C. Chique, *et al.*, "Cryptosporidium spp. in groundwater supplies intended for human consumption – A descriptive review of global prevalence, risk factors and knowledge gaps," *Water Research*, vol. 176. 2020. ISSN 0043-1354, <https://doi.org/10.1016/j.watres.2020.115726>.
- [15] F.Y. Ramírez-Castillo, *et al.*, "Waterborne pathogens: detection methods and challenges," *Pathogens*, vol. 4, no. 2, pp. 307–334, 2015. <https://doi.org/10.3390/pathogens4020307>.
- [16] P. Madoni. "Protozoa in wastewater treatment processes: A minireview," *Italian Journal of Zoology*, vol. 78, no. 1, pp. 3–11, 2011. <https://doi.org/10.1080/11250000903373797>.
- [17] K.K. Rosada, R. Adiarni N., and Sunardi, "Potensi Paramecium caudatum dalam mereduksi bahan organik dan bakteri coliform pada limbah cair rumah sakit," *Prosiding Seminar Nasional Jurusan Biologi FMIPA UNPAD*, pp. 314, 2013.
- [18] J.M. Baker, *et al.*, "The associations between water and sanitation and hookworm infection using cross-sectional data from Togo's national deworming program," *PLoS Negl Trop Dis*, vol. 12, no. 3. 2018. <https://doi.org/10.1371/journal.pntd.0006374>.