

## Case Report

# Cerebral Toxoplasmosis and Latent Tuberculosis Infection in HIV Infected Patient

Rijalun Arridho<sup>\*1</sup>, Raveinal<sup>2</sup>, Dwitya Elvira<sup>2</sup>

<sup>1</sup>Department of Internal Medicine, Faculty of Medicine, Universitas Andalas, Padang, 25175, Indonesia

<sup>2</sup>Division of Allergy and Immunology, Department of Internal Medicine, Faculty of Medicine, Universitas Andalas, Padang, 25175, Indonesia

\*Corresponding Author: [aldo.rijalun1996@gmail.com](mailto:aldo.rijalun1996@gmail.com)

### ARTICLE INFO

#### Article history:

Received 1 April 2024

Revised 30 March 2024

Accepted 19 April 2024

Available online 1 May 2024

E-ISSN: 2622-1357

P-ISSN: 2622-9234

#### How to cite:

Rijalun Arridho, Raveinal, Dwitya Elvira, "Cerebral Toxoplasmosis and Latent Tuberculosis Infection in HIV Infected Patient", Vol. 07, No. 02, May 2024.



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International. <https://doi.org/10.32734/sumej.v7i2.11843>

### ABSTRACT

**Background:** Toxoplasmosis is an opportunistic infection caused by *Toxoplasma gondii*, which affects one third of the global human population and commonly involves the central nervous system (CNS)/brain. The diagnosis of obvious sign of imaging method cerebral toxoplasmosis is commonly reported in several case report, but subclinical sign of imaging is uncommon reported. **Objective:** we reported a case of 28-years-old male with chief complaint headache that worsening since 3 days before admission. **Methods:** This is a case report. **Results:** He was first recognized as HIV infected patient on hospital admission. CT scan revealed minimal perifocal oedema in left frontal region. Patient was diagnosed as cerebral toxoplasmosis and latent tuberculosis infection in HIV clinical stadium IV. Patient received empirical therapy of cerebral toxoplasmosis and symptom relieved after 7 days of treatment. Cerebral toxoplasmosis typically associated with HIV infected patient that result in the reactivation of *Toxoplasma gondii* infection. **Conclusions:** Clinical presentation of CNS toxoplasmosis varies and normal brain CT can be found despite presumptive diagnosis was revealed.

**Keywords:** cerebral toxoplasmosis, empirical therapy, hiv infection, immunodeficiency

## 1. Introduction

Cigarette smoking is the main risk factor to cause lung cancer with 80-90% of estimated number. In a cigarette, around 4000 chemical substances, and some of them are carcinogenic matters, including polycyclic aromatic hydrocarbon, aromatic amines, nitrosamines, and organic/inorganic compounds such as benzene, vinyl chloride, arsenic and chromium. These carcinogenic matters have been reported to have relations to DNA mutations [1], and one of which is lung cancer that is triggered by the declining of immune system due to the blocking via smoking [2].

Human Immunodeficiency Virus (HIV) is a virus that can cause AIDS by attacking T lymphocytes especially CD4 which can cause decreased cellular immunity and increased occurrence of opportunistic infections, especially with CD4 T cell counts of less than 200 cells per mm<sup>3</sup>. One of opportunistic infection is cerebral toxoplasmosis with prevalence of cerebral toxoplasmosis is estimated approximately 30% of patients with AIDS. Although cerebral toxoplasmosis commonly present in AIDS, but subclinical abnormality of radiographic imaging is uncommon reported [1,2].

Cerebral toxoplasmosis or also called toxoplasmic encephalitis (TE) was one of opportunistic infection that was occurred. Prevalence of cerebral toxoplasmosis is estimated approximately 30% of patients with AIDS. in 56% of these subjects, the most common clinical manifestation was headache [2].

Cerebral toxoplasmosis is an infection of the brain caused by a parasite. The cause of toxoplasmosis disease is *Toxoplasma gondii* which is an obligate intracellular parasite [2]. Cerebral toxoplasmosis in immunodeficiency patients occurs due to reactivation of latent toxoplasma gondii infection. The prevalence of


cerebral toxoplasmosis in AIDS patients ranges from 3-40%. Patients with HIV infection who after an evaluation do not have active Tuberculosis (TB) should be treated as latent TB infection for at least 6 months [3].

## 2. Case Presentation

A 28-year-old male came with chief complaints of increased headaches since 3 days before admission. The patient presented with headache increased since 3 days ago which already felt since 2 weeks, continuously. Pain is felt in all parts of the head. Headache is not throbbing. Headache is not accompanied by vomiting. Decreased of appetite and weight loss since a month ago approximately 5 kg. Fever since 2 weeks ago. No history of chronic cough and night sweating. Patient was referred from Regional Hospital with reactive HIV result test and symptom headaches for further diagnosis and management. The patient status is homosexual and had sex history with his partner since 2018. Sex does not use condoms.

On physical examination, vital signs were obtained with temperature 38.3 C and VAS 4. General physical examination found within normal limits. Routine laboratory examinations within normal limits, ECG and X-ray thorax examinations also within normal limits. HIV confirmation test found positive on HIV confirmation test I, II, and III. HBsAg and anti-HCV lab test was non-reactive. Gene Xpert examination of sputum was negative for Mycobacterium Tuberculosis. We performed toxoplasma serology testing which shown IgG anti-toxoplasma positive with value 6.7 IU/mL (normal: < 1.6 IU/mL) and IgM anti-toxoplasma negative. Head CT scan non contrast showed that there is minimal focal edema in the left frontal region.

We diagnosed this patient as HIV clinical stage IV with cerebral toxoplasmosis and laten tuberculosis infection. The patient was given empirical therapy of cerebral toxoplasmosis, pyrimethamine oral loading dose 200 mg for 1 day and continue with 3 x 25 mg orally and clindamycin 4 x 600 mg. Antiretroviral treatment on this patient was FDC TLD (tenofovir-lamivudin-dolutegravir 2 x50 mg). we performed brain edema therapy given steroid injection of dexamethasone 4 x 5 mg IV, tapering off every 3 days. Latent TB infection therapy given isoniazid prophylaxis 1 x 300 mg PO. Preventive therapy for pneumocytis jirovecii pneumonia given cotrimoxazole 2 x 960 mg PO. After 1 week of empirical therapy of cerebral toxplasmosis, there was an improvement in headache symptoms.



**Figure 1.** (a) Axial section of CT scan, minimal perifocal oedema in left frontal region ; (b) chest x ray, normal.

## 3. Discussion

A 28-year-old man has been admitted to the internal medicine ward of Dr. M. Djamil Hospital on February 29, 2023 with a diagnosis HIV clinical stage IV with cerebral toxoplasmosis and latent tuberculosis infection. The diagnosis of obvious sign of imaging method cerebral toxoplasmosis is commonly reported in several case report, but subclinical sign of imaging is uncommon reported. The diagnosis of cerebral toxoplasmosis was established based on the clinical history of patients complaining of headaches since 2 weeks before hospital admission and increased since 3 days. The patient was referred from a regional hospital with HIV infection and severe headache. Head CT scan revealed that focal edema in the left frontal region. IgG anti-toxoplasma laboratory test was performed and positive results were obtained with value 6.7 IU/mL. Negative IgM anti-toxoplasma result was not excluded cerebral toxoplasmosis infection. Based on CDC guideline, IgG anti-

toxoplasmosis sufficient to positive serology result. Patient was given anti-toxoplasmosis therapy and showed a good response to treatment [4].

Headache is one of the most frequent clinical manifestations of toxoplasma infection in the central nervous system (CNS) of immunocompromised patients. Headache in cerebral toxoplasmosis patients can be caused by idiopathic intracranial hypertension, cerebral pseudotumor, and aseptic meningitis influenced by a wide variety of factors associated with indolamine 2,3-dioxygenase (IDO) and nitrite oxide (NO) [5].

In these patients, CT scan examination does not show a specific lesion in the brain, but only a focal edema in the left frontal region. Based on the diagnosis algorithm, if there is no mass in the brain, it is recommended to do a repeat examination. The main imaging examination in patients suspected of cerebral toxoplasmosis is head MRI with contrast. However, patients only performed head CT scan in order to lack of emergency diagnostic services.

Based on Graham, 2020, in some cases, especially asymptomatic toxoplasmosis, there were no abnormalities found on the CT scan of the head, but at autopsy found that there are multifocal cysts. MRI examination is recommended to this patient, because MRI is more sensitive to detect brain lesions. In these patients, headache symptoms improve after empirical therapy of cerebral toxoplasmosis so that a presumptive diagnosis can be established without further examination with MRI [6, 7].

Patients are given empirical therapy of cerebral toxoplasmosis, namely pyrimethamine and clindamycin. After clinical evaluation, symptoms improvement is obtained so that the presumptive diagnosis of cerebral toxoplasmosis can be established [8]. Presumptive therapy of toxoplasmosis encephalitis with pyrimethamine and clindamycin results are quite good. In more than 50% of cases within 3 days there is improvement. When therapy is given 7 days, the percentage that improves becomes higher (90%). Toxoplasmosis Encephalitis (TE)-related mortality in HIV patients is almost 100% if there is neglect of therapy [6, 8, 9].

The diagnosis of HIV stage IV clinical in this patient is established based on the examination of HIV confirmation tests was positive in I, II, and III tests. The patient has a history of fever since 2 weeks ago and claimed to have a habit of homosexual and having a sex with his partner since 2018. Laboratory test for CD4 revealed that decreased amount of cells 135 cells / uL.

Cerebral toxoplasmosis in immunodeficient patients occurs due to reactivation of latent toxoplasma gondii infection. The prevalence of cerebral toxoplasmosis in AIDS patients ranges from 3-40%. The clinical presentation of the patient can vary from fever, headache, sensory disturbances, motor impairment, focal neurological deficits, seizures, to loss of consciousness [6, 10].

In patients diagnosed as latent tuberculosis based on history and physical examination no signs and symptoms of tuberculosis infection were obtained, on TCM examination no Mycobacterium Tuberculosis was detected. According to the ISTC guideline, patients with HIV infection who after careful evaluation do not have active TB should be treated as latent TB infection given isoniazid 300 mg daily dose for at least 6 months [4, 11].

#### **4. Conclusion**

In conclusion, this case highlights the importance of recognizing cerebral toxoplasmosis in immunocompromised patients, particularly those with HIV infection. Despite the absence of specific lesions on head CT scan, the diagnosis was supported by positive serology, clinical symptoms, and good response to empirical therapy with pyrimethamine and clindamycin. Early detection and prompt treatment are critical, as delayed therapy significantly increases mortality. MRI remains the recommended imaging modality for better sensitivity, although clinical improvement following treatment can support a presumptive diagnosis. Management of latent tuberculosis infection is also essential in the comprehensive care of HIV patients to prevent future complications.

#### **5. Data Availability Statement**

The datasets generated and analyzed during the current study are not publicly available due to privacy and ethical considerations but are available from the corresponding author upon reasonable request.

#### **6. Ethical Statement**

Sumatera Medical Journal (SUMEJ) is a peer-reviewed electronic international journal. This statement clarifies ethical behavior of all parties involved in the act of publishing an article in Sumatera Medical Journal (SUMEJ), including the authors, the chief editor, the Editorial Board, the peer-reviewer and the publisher (TALENTA Publisher Universitas Sumatera Utara). This statement is based on COPE's Best Practice Guidelines for Journal Editors.

## 7. Author Contributions

All authors contributed to the design and implementation of this case report, data analysis, and finalizing the manuscript.

## 8. Funding

No funding.

## 9. Acknowledgements

The authors thank to the staff and member for their valuable assistance during data collection and analysis.

## 10. Conflict of Interest

Authors declares no conflict of interest.

## References

- [1] Fauci AS, Lane HC. Human immunodeficiency virus disease: AIDS and related disorders. In: Kasper DL, Fauci AS, Longo DL, Braunwald E, Hase SL, Jameson JL, editors. *Harrison's Principles of Internal Medicine*. 17th ed. New York: McGraw-Hill; 2010. p. 115-8.
- [2] Zawadzki R, Modzelewski S, Naumowicz M, Matyja AM, Urbaniak AD, Zajkowska J, et al. Evaluation of imaging methods in cerebral toxoplasmosis. *Pol J Radiol*. 2023;88:e389-98.
- [3] Vial JE. HIV-related cerebral toxoplasmosis revisited: current concepts and controversies of an old disease. *J Int Assoc Provid AIDS Care*. 2019;18:1-20.
- [4] Prandota J. Recurrent headache as the main symptom of acquired cerebral toxoplasmosis in nonhuman immunodeficiency virus infected subjects with no lymphadenopathy. *Am J Ther*. 2007;14:63-106.
- [5] Graham AK, Fong C, Naqvi A, Giang J. Toxoplasmosis of the central nervous system: manifestations vary with immune response. *J Neurol Sci*. 2020;11:1-13.
- [6] Harrison WT, Hulette C. Cerebral toxoplasmosis: a case report with correlation of radiographic imaging, surgical pathology, and autopsy findings. 2017;495-501.
- [7] Vidal JE. HIV-related cerebral toxoplasmosis revisited: current concepts and controversies of an old disease. *J Int Assoc Provid AIDS Care*. 2019;18:1-20.
- [8] Pawl T, Chaisson RE, Hopewell PC. Latent tuberculosis infection and HIV: advances and challenges in diagnosis and treatment. *Lancet Infect Dis*. 2019;19(3):e113-22.
- [9] Kelly J, Yousuf T, Mwaba P, Zumla A. Cerebral tuberculosis and cerebral toxoplasmosis in HIV-infected patients: clinical presentation, diagnosis, and management. *Lancet Neurol*. 2009;8(12):1150-8.
- [10] World Health Organization. *Latent tuberculosis infection: updated and consolidated guidelines for programmatic management*. Geneva: WHO Press; 2018.
- [11] Porter SB, Sande MA. Toxoplasmosis of the central nervous system in the acquired immunodeficiency syndrome. *N Engl J Med*. 1992;327(23):1643-8.