




SUMEJ

Sumatera Medical Journal

Journal homepage: <https://talenta.usu.ac.id/smj>



COVID-19 Pandemic Effect on Stress Levels and Sleep Patterns

Dirga Filannira Desky¹, Eka Roina Megawati^{*2}, Christoforos Giannaki³

¹Undergraduate Program of Medicine, Faculty of Medicine, Universitas Sumatera Utara, Medan 20155, Indonesia

²Department of Physiology, Faculty of Medicine, Universitas Sumatera Utara, Medan 20155, Indonesia

³Faculty of Life Sciences, University of Nicosia, Nicosia 2417, Cyprus

*Corresponding Author: eka3@usu.ac.id

ARTICLE INFO

Article history:

Received 28 June 2024

Revised 13 August 2024

Accepted 29 August 2024

Available online 5 September 2024

E-ISSN: 2622-1357

P-ISSN: 2622-9234

How to cite:

D.F. Desky, E.R. Megawati, C. Giannaki, "COVID-19 Pandemic Effect on Stress Levels and Sleep Patterns," *Sumatera Medical Journal*, vol. 07, no. 03, September 2024

ABSTRACT

Coronavirus disease (COVID-19) is an respiratory disease caused by the SARS-CoV-2 virus that cause serious illness and death. Physical distancing has important role to prevent and slow down its transmission but has some negative psychological effects such as stress, confusion, frustation, and sleep pattern changes. The aim of this study was to determine the effect of the COVID-19 pandemic on stress levels and sleep patterns. The subjects of this study were 100 medical students of Universitas Sumatera Utara. The data of this study was obtained by questionnaires and analyzed using Spearman correlation test. This study found that based on the stress level, there were 59 students categorized normal, mild 16 students, moderate 10 students, severe 9 students, and very severe 6 students. The sleep patterns described that only 11 students slept normally and 89 students experienced sleep disturbances during the COVID-19 pandemic. This study found there was a significant correlation between the COVID-19 pandemic on stress levels ($p:0.01$ and $r:0.58$) and sleep patterns ($p:0.01$ and $r:0.36$).

Keyword: COVID-19 pandemic, Sleep Patterns, Physical distancing, Stress level, Students



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International.

<http://doi.org/10.32734/sumej.v7i3.17080>

1. Introduction

Mental health during the COVID-19 pandemic is a serious matter, especially for students.[1], [2] Coronavirus disease 2019 (COVID-19) is an infectious disease with high transmissions like Middle East Respiratory Syndrome (MERS-CoV) and Severe Acute Respiratory Syndrome (SARS-CoV). SARS CoV2 is a new type virus that never been identified in humans, and on January 30, 2020, World Health Organization (WHO) has declared it as a Public Health Emergency of International Concern (PHEIC) [3].

According to the WHO in 2020, the COVID-19 pandemic had an impact on psychological status [4]-[5]. In order to stop the virus transmission, some policies have been made such as always wearing a mask, physical distancing, washing hands frequently, working from home (WFH), coughing and sneezing safely [6]. These policies will decrease people interaction among others that contribute to stress condition. Also the information about the fatal cause of COVID-19 that can be found anywhere has a role in leading to depression [7].

The Chinese survey showed that during the COVID-19 pandemic 53.8% of respondents experienced of moderate or severe symptoms psychological impact; 16.5% had moderate to severe depressive symptoms; 28.8% experienced moderate to severe anxiety symptoms, and 8.1% reported experiencing moderate to severe stress that would impact daily activities such as sleep patterns [7]. The psychological condition experienced by people is anxiety when infected, uncertainty about disease, and inability to adapt [8].

Another study in Egypt "Impact of the COVID-19 Pandemic on Mental Health and Social Support among Adult Egyptians" of 1,867 respondents also said 41.4% experienced a severe impact, 34.1% stress due to work, 55.7% financial stress, 62.7% were stressed due to being laid off, 53.9% experienced fear, 52% experienced feelings of helplessness and 66.3% felt worried [9]. When facing a COVID-19 pandemic situation, individuals will feel uncomfortable and less calm, which affects the quality of their sleep. Social support is needed such as a sense of empathy, accompanying individuals when they are experiencing problems, and creating a warm and comfortable atmosphere, making them feel cared for and loved so that individuals will be better to deal with problems [10].

Stress itself will result in psychological disturbances and activities related to the behavior undertaken.[6] During sleep well, melatonin is produced and increased that can reduce anxiety and results in a good immune system. The aim of this study was to examine the correlation between COVID-19 pandemic on stress levels and sleep patterns.

2. Method

This is an analytical study with a cross sectional design and the population of this study was students from the Faculty of Medicine, Universitas Sumatera Utara (USU). This study was conducted from June 2020 to January 2021. The sample was taken by stratified random sampling, the exclusion criteria for this study were students who consumed sedative drugs, certain substances that had insomnia effect, had or are being diagnosed by a doctor with severe mental disorders, and had a history of sleep disorders (insomnia).

This study was approved by Research Ethics Committee of Universitas Sumatera Utara with number 310/KEP/USU/2020. The pandemic situation was assessed by The COVID-19 pandemic questionnaire that had been validated previously, the stress level was determined by the Depression Anxiety Stress Scale [10] and the sleep patterns using the Pittsburgh Sleep Quality questionnaire.[11] Those questionnaires were spread by online using the social media as Google form link. The questionnaires of depression anxiety stress scale (references no 10) and the questionnaires of Pittsburgh sleep quality index (reference no 11) attached in this journal and the questionnaires had been translated into bahasa.

3. Result and Discussion

The characteristics of the subject were shown on Table 1. Most of the students had normal (59%) stress level and 89 students (89%) had sleep patterns disorder (see Table 1).

Table 1. Subjects Characteristics

Characteristics	n (%)
Sex	
Male	39
Female	61
COVID-19 pandemic	
High	78
Low	22
Stress level	
Normal	59
Mild	16
Moderate	10
Severe	9
Very severe	6
Sleep patterns	
Good	11
Poor	89

This study found there was a significant positive correlation between COVID-19 pandemic on stress levels (p value 0.01 and r value 0.58) and sleep patterns (p value 0.01 and r value 0.36) (Table 2).

Table 2. The correlations between COVID-19 pandemic on stress level and sleep patterns

Variables	COVID-19 pandemic		p value	r value
	Low	High		
Stress level				
Normal	32	27	0.01*	0.58

Mild	0	16		
Moderate	0	10		
Severe	1	8		
Very severe	0	6		
Sleep patterns				
Good	10	1	0.01*	0.36
Poor	70	19		

*p value<0.05, Spearman correlation test

Discussions

This study found there was a significant correlation between COVID-19 pandemic on stress level and sleep patterns. This indicated that the COVID-19 pandemic is still a difficult thing to accept in every circle, especially medical students due to changes in the learning system from face to face to the online system [12]. Previous study on 7.143 students showed that 0.9% of students had severe anxiety, 2.7% had moderate anxiety, and 21.3% had mild anxiety. In addition, having relatives or acquaintances infected with COVID-19 is a risk factor for student of becoming anxiety (OR = 3.007, 95% CI = 2.377 – 3.804) [13]. Stress itself focuses on coping mechanisms when responding problems [14]. Most of the stress on students caused by a large amount of material to be studied, lack of free time, and academic pressure that is full of competition and separation from loved ones, also decreases productivity [15]. The status of pandemic or the global epidemic indicated that the transmission of COVID-19 was very fast and this status forced the government to implement some new policies such as work from home and social distancing [16]. This condition came suddenly making people unprepared to face it physically and psychologically [17]. Among the psychological conditions experienced by society is the feeling of anxiety about being infected [18]. Anxiety is an emotional state that arises when an individual is stressed, and is characterized by tense feelings, worried and accompanied by physical responses such as a fast heartbeat and increased blood pressure [19].

Another study on medical students in 2016 at Universitas Sumatera Utara with chi-square test analysis showed that there was a significant correlation between stress levels and sleep quality with values ($r = 0.595$ and $p = 0.001$), [20] that indicated the higher the stress level, the worse the sleep quality [21]. Chronic insomnia also deteriorates and acute insomnia is now popping up because the difficulty limits the rest time and work time during the isolation period at home so that it is difficult to fall asleep when you have to sleep which adversely affects sleep continuity to lack of sleep at night and disturbing activity during the day [22].

A web-based study totaling 7.236 volunteers with a cross-sectional method and multivariate analysis test found 35.1% generalized anxiety disorder (GAD), 20.1% depressive symptoms, and 18.2% sleep quality disorders. It shows that health care workers are at high risk of poor sleep quality, also the prevalence of GAD <35 years in younger people, depressive symptoms is much higher than in older people [23].

4. Conclusion

This study found that the COVID-19 pandemic is closely related to both psychological and mental health, one of which also has impact on stress level and sleep patterns, the inability to adapt in the middle of the pandemic will cause anxiety, as well as the ambiguity of disease and limited activities to make this sustainable.

Ethics Approval: Sumatera Medical Journal (SUMEJ) is a peer-reviewed electronic international journal. This statement below clarifies ethical behavior of all parties involved in the act of publishing an article in Sumatera Medical Journal (SUMEJ), including the authors, the chief editor, the Editorial Board, the peer-reviewer and the publisher (TALENTA Publisher Universitas Sumatera Utara). This statement is based on COPE's Best Practice Guidelines for Journal Editors.

Authors Contributions: All authors contributed to the design and implementation of the research, data analysis, and finalizing the manuscript.

Funding: No funding.

Disclosure: Authors declares no conflict of interest.

References

- [1] R. Shrestha, L. Shrestha, *et al.*, "Coronavirus disease 2019 (Covid-19): A pediatric perspective," *J. Nepal Med. Assoc.*, vol. 58, no. 227, pp. 525–532, 2020. <https://doi.org/10.31729/jnma.4977>.
- [2] E.J. Emanuel, *et al.*, "Fair Allocation of Scarce Medical Resources in the Time of Covid-19," *N. Engl. J. Med.*, vol. 382, no. 21, pp. 2049–2055, 2020. <https://doi.org/10.1056/nejmsb2005114>.

- [3] WHO, "WHO/Europe | Coronavirus disease (COVID-19) outbreak - Mental health and psychological resilience during the COVID-19 pandemic," 2020.
- [4] T. Respati, H.S. Rathomi, "Bunga Rampai Artikel Penyakit Virus Korona (COVID-19)." Pusat Penerbitan UNISBA (P2U), 2020.
- [5] Y. Shu, H. He, X. Shi, Y. Lei, and J. Li, "Coronavirus disease-2019 (Review)," *World Acad. Sci. J.*, vol. 3, no. 2, pp. 1–7, 2021. <https://doi.org/10.3892/wasj.2021.83>.
- [6] C.S. Ho, C.Y. Chee, and R.C. Ho, "Mental Health Strategies to Combat the Psychological Impact of COVID-19 Beyond Paranoia and Panic," *Ann. Acad. Med. Singapore*, vol. 49, no. 1, pp. 1–3, 2020.
- [7] S.K. Brooks, *et al.*, "The psychological impact of quarantine and how to reduce it: rapid review of the evidence," *Lancet*, vol. 395, no. 10227, pp. 912–920, 2020. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8).
- [8] S.M. El-Zoghby, E.M. Soltan, and H.M. Salama, "Impact of the COVID-19 Pandemic on Mental Health and Social Support among Adult Egyptians," *J. Community Health*, vol. 45, no. 4, pp. 689–695, 2020. <https://doi.org/10.1007/s10900-020-00853-5>.
- [9] F.P.J.D.W. Johnson, *Joining together: Group theory and group skills.*, vol. 21, no. 10. 2012. <https://doi.org/10.1037/014685>.
- [10] H. Lovibond, S and F. Lovibond, P, "Depression Anxiety Stress Scale (DASS) 2021," *Psychology*, pp. 20–22, 2018.
- [11] D.J. Buysse, C.F. Reynolds, T.H. Monk, S.R. Berman, and D.J. Kupfer, "The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research," *Psychiatry Res*, vol. 28, pp. 193–213, 1989.
- [12] W. Cao, *et al.*, "The psychological impact of the COVID-19 epidemic on college students in China," *Psychiatry Res.*, vol. 287, pp. 112934, 2020. <https://doi.org/10.1016/j.psychres.2020.112934>.
- [13] R.D. Rahmayani, R.G. Liza, and N.A. Syah, "Gambaran Tingkat Stres Berdasarkan Stressor pada Mahasiswa Kedokteran Tahun Pertama Program Studi Profesi Dokter Fakultas Kedokteran Universitas Andalas Angkatan 2017," *J. Kesehat. Andalas*, vol. 8, no. 1, pp. 103, 2019, <https://doi.org/10.25077/jka.v8i1.977>.
- [14] Y. Livana, Mubin, & Basthomi, "Penyebab Stres Mahasiswa Selama Pandemi Covid-19," *Jurnal Ilmu Keperawatan Jiwa*, vol. 3, no. 2, pp. 203–208, 2020.
- [15] S. dwi Putri, "Hubungan Tingkat Stres Dengan Kualitas Tidur Pada Mahasiswa Semester VII Fakultas Kedokteran Universitas Sumatera Utara Tahun 2016," *Skripsi*, pp. 91, 2016.
- [16] T.A, "COVID-19 dan lansia," *Pus. Pnb. Unisba LPPM UNISBA*, 2020.
- [17] S.A and M. Phil, "Gambaran umum persepsi masyarakat terhadap bencana di Indonesia," *J. Ilmu Ekon. dan Sos.*, vol. 5, no. 3, pp. 304–326, 2016.
- [18] L. Fitria and I. Ifdil, "Kecemasan remaja pada masa pandemi Covid-19," *J. Educ. (Jurnal Pendidik. Indones.)*, vol. 6, no. 1, pp. 1–4, 2020. <https://doi.org/10.37287/jppp.v3i3.530>.
- [19] S.A. Beaudreau and R. O'Hara, "The Association of Anxiety and Depressive Symptoms With Cognitive Performance in Community-Dwelling Older Adults," *Psychol. Aging*, vol. 24, no. 2, pp. 507–512, 2009. <https://doi.org/10.1037/a0016035>.
- [20] H. Xiao, Y. Zhang, D. Kong, S. Li, and N. Yang, "Social capital and sleep quality in individuals who self-isolated for 14 days during the coronavirus disease 2019 (COVID-19) outbreak in January 2020 in China," *Med. Sci. Monit.*, vol. 26, pp. 1–8, 2020. <https://doi.org/10.12659/MSM.923921>.
- [21] D. Yulia, *et al.*, "Insomnia During COVID-19 Pandemic," *J. Ilm. Kesehat. Sandi Husada*, vol. 9, no. 2, pp. 1111–1116, 2020. <https://doi.org/10.35816/jiskh.v10i2.483>.
- [22] N. Vindegaard and M.E. Benros, "COVID-19 pandemic and mental health consequences: Systematic review of the current evidence," *Brain. Behav. Immun.*, vol. 89, pp. 531–542, 2020. <https://doi.org/10.1016/j.bbi.2020.05.048>.
- [23] Y. Huang and N. Zhao, "Corrigendum to Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: a web-based cross-sectional survey [Psychiatry Research, 288 (2020) 112954]," *Psychiatry Res.*, vol. 299, pp. 113803, 2021. <https://doi.org/10.1016/j.psychres.2021.113803>.