

The Comparison of Er α Expression Between Endometriosis with No Endometriosis

D. Irsat Syafardi^{1*}, **Henry Salim Siregar**², **Yostoto B Kaban**³, **Edy Ardiansyah**⁴, **Riza Rivany**⁵

^{1,2,3,4,5} Department of Obstetrics Gynecology, Faculty of Medicine Universitas Sumatera Utara, Indonesia

Abstract. Background: Estrogen receptors (ER) play an important role in mediating action. The ER α has a higher affinity for estrogen and the dominant form of the normal endometrium. A cross-sectional study from an ectopic tissue of endometriosis and normal was examined for immunohistochemistry. This research was conducted from November 2015 until the sample complete. The analysis was performed using Fisher Exact test, $p < 0.05$ was considered to any difference of estrogen alpha receptor expression between endometriosis patients with no endometriosis. RE α is more dominant in normal endometrium, whereas in the case of endometriosis in the presence of many negative expression of RE α then the assumption that RE β is a receptor that many encountered. There is a difference in the expression of RE α between endometriosis with no endometriosis.

Keyword: Estrogen A Receptor, Endometriosis, Normal Endometrium.

Abstrak. Reseptor estrogen (ER) memainkan peran penting dalam mediasi tindakan. ER α memiliki afinitas yang lebih tinggi untuk estrogen dan bentuk dominan dari endometrium normal. Sebuah studi cross-sectional dari jaringan ektopik endometriosis dan normal diperiksa untuk imunohistokimia. Penelitian ini dilakukan dari November 2015 hingga sampel selesai. Analisis dilakukan dengan menggunakan uji Fisher Exact, $p < 0,05$ dianggap sebagai perbedaan ekspresi reseptor alpha estrogen antara pasien endometriosis tanpa endometriosis. RE α lebih dominan pada endometrium normal, sedangkan pada kasus endometriosis di hadapan banyak ekspresi negatif RE α maka anggapan bahwa RE β adalah reseptor yang banyak ditemui. Ada perbedaan dalam ekspresi RE α antara endometriosis tanpa endometriosis.

Kata Kunci: Reseptor Estrogen A, Endometriosis, Endometrium Normal.

Received 31 December 2018 | Revised 17 January 2019 | Accepted 31 January 2019

1. Introduction

Endometriosis is defined as the appearance of endometrial tissue outside of the uterine cavity. Endometriosis is strongly influenced by estrogen measure. In the United States, this disease affects 5 - 19% of women at the reproductive age. The main descriptions are chronic pelvic pain

*Corresponding author at: Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia

during coital and infertility [1]. Pelvic pain and complaints from endometriosis can be reduced by giving the anti-inflammatory drugs, so it supports that in endometriosis, the process of inflammatory gives the contribution as one of the pathogenesis of the disease [2].

The Proliferation of endometrial cells and endometriosis is induced by estrogen. In contrast, the progesterone stimulates cellular differentiation and suppresses the cellular proliferation. In the endometriosis, the increasing estrogen effects and abnormal progesterone work, leading to the increasing cell proliferation. Simultaneously, apoptotic disorders in the endometrial and endometrial cells from women with endometriosis can contribute to the pathogenesis of the disease [3].

The Estrogen receptor has an important role in mediating estrogen action in tissues. There are two forms of estrogen receptors, they are ER α and ER β , which are encoded by different genes. The estrogen α receptor has a higher affinity for estrogen, and it is the dominant form of the normal endometrium. Recent studies have shown that ER α and ER β express the human tissue endometriosis, but the distribution of iso form is different between eutopic endometrium and ovarian, endometrial tissue [4], [5].

From the description that has been described above, the researchers are very interested to know whether there is difference expression of RE α on ectopic endometrial tissue endometriosis patients and normal endometrial tissue to further know about endometriosis disease.

2. Methodology

This was an analytic study with cross-sectional design in which it has been done the examination of immune histo hemical toward the ectopic endometrial endophilic tissue blocks of endometriosis and normal endometrial tissue. The study was conducted at Department of Obstetrics and Gynecology FK USU / RSUP H Adam Malik Medan, while immune histochemical examination was done at Department of Anatomical Pathology USU Medan. This research was conducted from November 2015 until the sample was met. This research had been approved by the Committee of Ethics of the Faculty of Medicine, University of Sumatera Utara. The preparations cannot be analyzed by the cause of poor paraffin preparation removed from the study. The immune histochemical assessment for the expression of the RE antibody uses Allred's score by summing up the percentage of the colored cell / Proportion Score (PS) and the intensity score of the staining / Intensity Score (IS). It is said negative expression if the score shows 0-2, while positive if the score shows ≥ 3 .

2.1 Statistical analysis

The analysis of the research was conducted by using Fisher Exact test, $p < 0.05$ was considered the different of the expression of RE α between endometriosis patients with no endometriosis.

3. Results

This study has been done by using the subjects of cases of endometriosis as many as 23 people and control of non-Endometriosis (normal endometrium) as many as 23 people. The following are the characteristics of the research subjects we obtained (**Table 1**).

Table 1 Frequency Distribution Based on the Characteristics of research subjects based on Age and Parity

Characteristic	Research Group			
	Endometriosis		Normal	
	n	%	n	%
Age				
< 30	11	47.9	11	47.9
30 – 40	8	34.8	12	52.1
> 40	4	17.3	0	0
Total	23	100.0	23	100.0
Parity				
Nullipara	23	100.0	7	30.4
Primipara	0	0	7	30.4
Multipara	0	0	9	39.2
Total	23	100.0	23	100.0

From the results of research, we also found the distribution of research subjects based on the endometriosis stage as seen in **Table 2** below.

Table 2 Frequency Distribution of Respondents Based on Endometriosis Stage

Stage	Endometriosis	
	n	%
Stage I (minimum)	0	0
Stage II (light)	2	8.7
Stage III (medium)	10	43.5
Stage IV (weight)	11	47.8
Total	23	100

In this study, it's found that there was a significant difference in the expression of alpha estrogen receptor on endometriosis group stroma with non-endometriosis, and the expression value of alpha estrogen receptor in the epithelial group of endometriosis with non-endometriosis as seen in **Table 3** and four below.

Table 3 Differentiation of Alfa Estrogen Receptor Expression on Stroma Group Endometriosis and Non-Endometriosis

Expression of Estrogen Receptor α (Stroma)	Research Group		<i>p</i>
	Endometriosis	Normal	
Negative (Score 0-2)	21 (91.3%)	9(39.2%)	.000*
Positive (Score \geq 3)	2 (8.7%)	14(60.8%)	

*Test Fisher exact

Table 4 Differences Expression of Alfa Estrogen Receptor on Epithelial Endometriosis and Non-Endometriosis

Expression of Estrogen Receptor α (Epitel)	Research Group		<i>p</i>
	Endometriosis	Normal	
Negative (Score 0 - 2)	23 (100%)	11 (47.8%)	.000*
Positive (Score \geq 3)	0 (0%)	12 (52.2%)	

*Test Fisher exact

The Endometriosis occurrences are rarely found in the teenage girl of primers, but it can be identified in more than 50% of women younger than 20 years with chronic pelvic pain or dyspareunia. The average age in which a patient is diagnosed with endometriosis is between the ages of 25-30 years [4]–[6]. In this study, the largest group of endometriosis in under 30 years was 11 people (47.9%), while the whole patient was nullipara (100%).

Endometriosis is an inflammatory disease that depends on estrogen levels, due to P450 aromatase and deficiency of 17 Beta-hydroxysteroid de-hydrogenase. 17 Beta-hydroxysteroid dehydrogenase converts estradiol into a less active estrogen, which is not found in the luteal phase of endometrial tissue. Therefore, endometriosis occurs in many women at the reproductive age [2], [7].

The exact incidence of endometriosis in Indonesia currently has not known yet for sure. At least 20-40% of women with infertility have endometriosis. Meiling H stated that high parity would reduce the risk of endometriosis [4], [8], [9].

Endometriosis tissue has a high rate of proliferation. Proliferation is associated with the progression of disease that characterized by the ability to form large masses such as endometriosis cysts and invasion of adjacent tissues that characterized by degrees of disease or staging. In this study, we encountered cases of endometriosis mostly at an advanced stage. We also found a significant difference in the value of alpha estrogen receptor expression in the endometriosis group stroma with non-endometriosis, and the expression value of alpha estrogen receptors in epithelial endometriosis with non-Endometriosis.

States that alpha estrogen receptor expression in stromal endometriosis cells is lower when compared to the normal endometrial stroma. This is most likely due to the failed response of estradiol (E2) to induce expression of progesterone (PR), which ultimately leads to secondary progesterone deficiency and progesterone resistance in the study subjects. The alpha-estrogen receptor gene in humans itself is regulated by various types of promoters [8], [10]. Says that stromal cells can produce the paracrine growth factors necessary to induce the proliferation of epithelial cells by estrogen. RE- α and RE- β are proteins with high affinity for E2 and encoded by separate genes. Although RE- α and RE- β are present in the endometrium, RE- α is the major mediator of estrogenic action in endometrial tissue.

The human endometrium is a unique tissue that can cycle in proliferation, differentiation, implantation, and regeneration in response to ovarian steroid production. The effects of estrogen itself are mediated by two different estrogen receptors, RE α and RE β , which are families of nuclear receptors and act as activating transcription factors. Biological action of estrogen in the target tissue is modulated by hormone levels and receptor distribution. In the normal endometrium both RE α and RE β are encountered, but with different expressions during the menstrual cycle, where the RE α is more expressed than RE β in the normal endometrium, therefore estrogen-mediated proliferation is largely due to activation of RE α [11].

Further research is needed on the ratio of estrogen and beta estrogen receptors to normal endometriosis and endometrium. Also, there is a need to take a deeper look at other biomolecules concerning the use of drugs that act as selective estrogens modulator to regulate estrogen levels that affect the occurrence of endometriosis.

4. Conclusions

In the normal endometrium, RE α is more dominant, whereas in the case of endometriosis in the presence of many negative expression of RE α then the assumption that RE β is a receptor that is mostly being encountered.

REFERENCES

- [1] J. . Lund, *The Estrogen Receptor*. University of Aarhus:1-43, 2005.
- [2] D. Xu *et al.*, "The wedelolactone derivative inhibits estrogen receptor-mediated breast, endometrial, and ovarian cancer cells growth," *BioMed research international*, vol. 2014, 2014.
- [3] S. C. Hewitt and K. S. Korach, "Estrogen receptors: structure, mechanisms and function," *Reviews in Endocrine and Metabolic Disorders*, vol. 3, no. 3, pp. 193–200, 2002.
- [4] K. Zulli, B. Bianco, F. A. Mafra, J. S. Teles, D. M. Christofolini, and C. P. Barbosa, "Polymorphism of the estrogen receptor β gene is related to infertility and infertility-associated endometriosis," *Arquivos Brasileiros de Endocrinologia & Metabologia*, vol. 54, no. 6, pp. 567–571, 2010.
- [5] K. Tamm, M. Suhorutshenko, M. Rõõm, J. Simm, and M. Metsis, "The tissue specific role of estrogen and progesterone in human endometrium and mammary gland," in *Steroids-Basic Science*, InTech, 2012.

- [6] S. E. Bulun *et al.*, “Estrogen receptor- β , estrogen receptor- α , and progesterone resistance in endometriosis,” in *Seminars in reproductive medicine*, 2010, vol. 28, p. 36.
- [7] Z. Weihua *et al.*, “Estrogen receptor (ER) β , a modulator of ER α in the uterus,” *Proceedings of the National Academy of Sciences*, vol. 97, no. 11, pp. 5936–5941, 2000.
- [8] S. E. Bulun, “Endometriosis,” *New England Journal of Medicine*, vol. 360, no. 3, pp. 268–279, 2009.
- [9] M. Han, L. Pan, B. Wu, and X. Bian, “A case-control epidemiologic study of endometriosis,” *Chinese medical sciences journal= Chung-kuo i hsueh k'o hsueh tsa chih*, vol. 9, no. 2, pp. 114–118, 1994.
- [10] E. Trukhacheva, Z. Lin, S. Reierstad, Y.-H. Cheng, M. Milad, and S. E. Bulun, “Estrogen receptor (ER) β regulates ER α expression in stromal cells derived from ovarian endometriosis,” *The Journal of Clinical Endocrinology & Metabolism*, vol. 94, no. 2, pp. 615–622, 2009.
- [11] C. Pellegrini *et al.*, “The expression of estrogen receptors as well as GREB1, c-MYC, and cyclin D1, estrogen-regulated genes implicated in proliferation, is increased in peritoneal endometriosis,” *Fertility and sterility*, vol. 98, no. 5, pp. 1200–1208, 2012.