

IDENTIFICATION OF CANDIDA SPECIES ON THE SKIN OF DIABETES MELLITUS PATIENTS

OK. Muhammad Al Hafiz¹, Rina Amelia², Nelva Karmila Jusuf³

¹Participant of Educational Doctor Profession Program, Faculty of Medicine Universitas Sumatera Utara, Indonesia

²Department of Medicine Community, Faculty of Medicine Universitas Sumatera Utara, Indonesia

³Department of Dermatovenerology, Faculty of Medicine Universitas Sumatera Utara, Indonesia

Abstract— Background: Intertriginous candidiasis is a type of candidiasis which the lesion lies in the folds of the armpit, groin, intergluteal, folding of the breast, between the fingers or the toes, glans penis, and the umbilicus. One of the risk factors for intertriginous candidiasis is diabetes mellitus (DM). DM is a typical clinical syndrome characterized by the presence of hyperglycemia which is caused by deficiency or the decrease of insulin effectivity. Fungal infection of the skin with *Candida* species becomes easier in patients with immunocompromised conditions such as DM patients.

Objective: To identify the *Candida* species in DM patient's skin.

Method: This research is an observational descriptive study with cross sectional design. This research identifies *Candida* species from 30 DM patients at The Public Health Center Sentosa Baru Medan who met the inclusion and exclusion criteria by skin scraping with KOH 10% and slide culture.

RESULTS: The location of the rash was found in the crotch folds 63.3% and in the intergluteal 36.7%. All samples (100%) showed positive fungus structure with KOH 10% examination. On the examination of slide culture with Saboraoud Dextrose Agar media and identification of slide colony with *Corn Meal Agar* medium as many as 60% *Candida* spesies was found with overall species are *Candida albicans*.

Conclusions: The identification of *Candida* Species on the skin of DM patients showed as many as 60% *Candida albicans* was found.

Keywords— *Candida* Species, Intertriginosa Candidiasis, Diabetes Mellitus (DM)

I. INTRODUCTION

Candida is a fungal consisting of many species, but only 17 species are reported to infect humans. These species include *Candida albicans*, *Candida glabrata*, *Candida parapsilosis*, *Candida tropicalis*, *Candida krusei*, *Candida kefyr*, *Candida guilliermondii*, *Candida lusitaniae*, *Candida dubliniensis*.¹⁻⁴

Candidiasis can be found all over the world, relatively common in the tropical area that can affect all ages, both male and female.^{5,6} Research conducted at *National Skin Center*, Singapore, from 1999-2003 placed candidiasis (11.1%) in the third most cases of mycosis after dermatophytosis (57%) and pityriasis versicolor (25.2%). From all of the studies were found, most of the cases are intertriginous candidiasis (69.8%). This is due to the hot and humid climate of Singapore.⁷ Similar results were found in studies conducted in Bangladesh, where candidiasis ranks the third positions of superficial fungal infections.⁸

Candida infection occurs when there are factors that can nourish the growth or facilitate the invasion of the tissues due to the weak resistance of the host. The factors that cause candidiasis are divided into two, endogenous and exogenous predisposing factors. Endogenous predisposing factors such as pregnancy (due to pH changes in the vagina), obesity (due to sweat), iatrogenic, endocrinopatya (DM), chronic disease (tuberculosis, lupus erythematosus with poor general circumstances), age (elderly and younger infants exposed to the infection due to imperfect immunological status), immunologic (genetic disease). Exogenous predisposing factors such as:

climate, skin hygiene, foot bath habit and contact with patients.⁵

One form of candidiasis cutis is intertriginous candidiasis. Intertriginous candidiasis lies in the area of the underarm skin fold, groin, intergluteal, folding of the breast, between the fingers or toes, glans penis, and umbilicus, sirkumskripta, scaly, wet, and erythematous. The lesion is surrounded by satellites such as tiny vesicles and pustules or bullae which, when it breaks out it leaves erosive areas, with rough edges and develops as primary lesions.⁵

As known that DM is a predisposing factor of intertriginous candidiasis caused by the changes in the body's defense system.⁹⁻¹⁴

DM is a hyperglycemia disease characterized by absolute absence of insulin or a relative decrease in cell insensitivity to insulin.¹⁵ The metabolic status of DM patients provides the advantage by fulfilling specific nutritional requirements and facilitating the growth of fungi, especially *Candida* species.¹⁶ In a study at the General Hospital of Dr. Soetomo Surabaya, DM as a risk factor for the occurrence of intertriginosa candidiasis. Where the average patients with history of DM had a 3.26 times greater risk than no history of DM.

The purpose of this study was to identify *Candida* species on the skin of DM patients.

II. METHODOLOGY

This research is an observational research. The studies used in this research is cross sectional. This study was conducted from March to December 2016 at the Sentosa Baru Public Health Center Medan because it has many DM patients so that fullfill the inclusion criteria where it has a description of skin disorders in the armpit, groin, intergluteal, fingers or toes, glans penis and umbilicus and still a DM patient at Sentosa Baru Health Center Medan. Data that have been collected at the time of filling patient status data then examined skin scrapping with KOH 10% inspection and fungal culture in the laboratory of Microbiology FK USU.

III. RESULTS

The research was conducted in Sentosa Baru Health Center working area located at Jalan Sentosa Baru

No. 22 Kecamatan Medan Perjuangan Medan, Indonesia. Total of samples in this research are 30 people.

TABEL 1
THE DISTRIBUTION OF DM PATIENTS BY SEX

Sex	N	%
Male	12	40
Female	18	60
Total	30	100

Table 1 from the table above it is found that the most patient were female with a total of 60% and the rast were men with a total of 40%. Based on the 2007 National Basic Health Research report in Indonesia (RISKENDAS 2007) DM is more common found in woman with a percentage of 64.9% and men with a percentage of 35.1%.¹⁸ Maulina et al in H.Adam Malik General Hospital Medan found that most of the patient were with percentage of female 61.9% while with percentage of gender of men 38.1%.¹⁹

TABEL 2.
THE DISTRIBUTION OF DM PATIENTS BY AGE

Age	n	%
46 – 55 (initial elderly)	5	16,7
56 – 65 (late age)	17	56,7
>65 (elderly)	8	26,7
Total	30	100

Table 2 The table above found most people was found to suffer more from skin disease in the 56-65 year age group of 56.7%. In the study, Maulina et al found that intertriginous candidiasis was most common in the age group 45-65 years (45.95%) and followed by age group 65 years (21.62%). The age group above 50 years ranks first with 40% and the age group 30 to 50 years with 15.5%.¹⁸ This is because immune system status in the elderly has decieased. In addition, elderly people are susceptible to some diseases such as diabetes mellitus, which is one of the risk factor of intertriginous candidiasis.²⁰

TABEL 3.
THE DISTRIBUTION OF DM PATIENTS BY THE LEVEL OF
EDUCATION

Education Level	n	%
Primary School	4	13,3
Junior High School	11	36,7
Senior High School	12	40
Bachelor	3	10
Total	30	100

Table 3 The table above found that the incidence of candidiasis 40% in high school students and followed by 36.7% of junior high school students. Knowledge of a person is very closely related to the achieved education. People who have low level of education, knowledge about health is certainly not too profound. This could be the reason of them not knowing DM disease in depth. Other behaviors are based on knowledge and positive attitude.²¹ The patient's knowledge of Diabetes Mellitus is a tool that helps the patient manage Diabetes during their lifetime. The more and better the patients understand their illness, the more they understand how to change their behavior and lifestyle, and why it is needed.²²

TABEL 4.
THE DISTRIBUTION OF DM PATIENTS BASED ON THE
DURATION IN SUFFERING IN DIABETES MELLITUS

The duration	n	%
<1 year	3	10
1 – 5 year	7	23,3
6 – 10 year	7	23,3
11 – 20 year	9	40
>20 year	4	13,3
Total	30	100

Table 4 from the table above it was found that the duration of suffering DM for 11-20 years was found in 9 people (30%). The long duration of DM affects patients' behavior toward treatment, patients who are still newly diagnosed with DM generally will be open and happy to be given counseling about drugs, because they are still not familiar with the disease and treatment experienced, so there is a great sense of curiosity towards the disease and its treatment. While patients who have

long suffered with DM, they will assume that the disease is not dangerous, or in their experience the results are not so satisfactory during their treatment, they will resigned and less concerned about the disease so they become less interested when given information about their diseases and medicines.²³

TABEL 5.
THE DISTRIBUTION OF DM PATIENTS BASED ON THE LOCATION
OF THE RASH

The Location of the Rash	n	%
Groins	19	63,3
Intergluteal	11	36,7
Total	30	100


Table 5 from the table above it was found that the rash in the skin fold areas affected by intertriginous candidiasis gave a result of 19 people (63.3%) in the groin area. This is because the groins are the most closed part when someone is dressed. When the temperature of the environment is hot and a person is sweating, the groins become the most moist part because of the sweat. This condition supports the growth of candida fungal in the groin and eventually leads to intertriginous candidiasis.²⁰

TABEL 6.
10% KOH EXAMINATION

KOH 10%	n	%
Hyphae (+) Spore(+)	2	6,7
Hyphae (+) Spore (-)	0	0
Hyphae(-) Spore (+)	28	93,3
Hyphae(-) Spore (-)	0	0
Total	30	100

Table 6 From the table above, it was found 10% KOH examination results in its function to dissolve debris and lipids from skin, hair, and mucous scrapings. 30 samples was tested with KOH 10% examination, where hyphae (-) and spore (+) was found in 28 samples (93.3%) and the hyphae (+) spotted (artroconidia) spore (-) in 2 samples (6.7%). Next, the fungal culture will be examined. Morphologically has several forms of yeast elements such as yeast cells (blastopora/yeast), hyphae and intermedia/pseudohifa form. The yeast cells are round, oval or round oval with size 2-5 μ x 5-28 μ . Candida multiplies itself by forming bud

that will continue to elongate to form a pseudohyphae.²⁴



A.

A. Hyphae (-) Spore (+)

B.

B. Hyphae (+) Spore (+)

TABEL 7.
OVERVIEW OF FUNGAL CULTURE

Yeast species	n	%
<i>Candida albicans</i>	18	60
No <i>Candida</i> was found	12	40
Total	30	100

Table 7 From the table above, it shows that 18 samples of *Candida albicans* (60%) and no *Candida* was found in 12 samples (40%). This is in line with Maulina's and her friends' findings. Where they found that *Candida albicans* as much as 67%.¹⁹



A. Chromogenik Medium

B. Slide Culture

Identification of species can be done macroscopically and microscopically. Macroscopically it can be done on chromogenic media (CHROMagar). In this medium *Candida* species will form different colony colors. *Candida albicans* form a green colony.²⁵

Identification of microscopic morphologic species can be done by fungal culture on certain mediums, such as *Corn Meal Agar*. In this medium *Candida albicans* forms a terminal klamidospora

which is a large yeast cell with thick walls and is located at the end of the hyphae.²⁴

One of the risk factors in intertriginous candidiasis is DM. The condition of chronic hyperglycemia in diabetic patients is associated with long-term complications that one of them attacks the skin.²⁶ The skin becomes one of the organs often affected by DM. Skin manifestations of infection become one of the most common chronic complications seen in DM patients.²⁷

In addition, there are several other predisposing factors found in the respondents when they were filling the status of DM participants where there are areas of moist skin folds and lack of maintaining the cleanliness of the skin area. From the daily bathing habits, the use of soap which is only occasionally, the frequency of changing clothes in a day it was found only 1 time or 2 times in a day, where the lack of behavior to keep the skin area clean and sweaty habits that found moisture in the skin folds. This condition supports the growth of *Candida* fungus in the skin folds and eventually leads to intertriginous candidiasis.

IV. CONCLUSIONS

The identification of *Candida* species in DM patient skin found that there were *Candida albicans* species as much as 60%.

SUGGESTION

This study can be continued by identification of other fungi such as dermatophytes in DM patients. In the need for counseling and plenary services for DM patients and skin disorders associated with it.

REFERENCES

- [1] Verma S, Heffernan MP. Superficial fungal infection. In: Wolf K, Goldsmith LA, Katz SI, Gilchesrt BA, Paller AS, Leffell DJ, editors. Fitzpatrick's Dermatology In General Medicine. 7thed. New York: McGraw-Hill Companies Inc, 2008.p.18077-44
- [2] Aliyu SH, Enoch DA, Abubakar II, Ali R, Carmichael AJ, Farrington M, et al. Candidaemia in large teaching hospital: a clinical audit. Q J Med. 2006;99:655-63.
- [3] Wahyuningsih R, Freisleben HJ, Sonntag HG, Schnitzler P. Simple and rapid detection of *C. albicans* DNA in serum by PCR for diagnosis of invasive candidiasis. J Clin Microbiol. 2000;38:3016-21.
- [4] Dismukes WE, Pappas PG, Sobel JD. Clinical Mycology. Oxford University Press America. 2003.
- [5] Djuanda A, Hamzah M, Aisah S, editor. Ilmu Penyakit Kulit dan Kelamin. Edisi ke-6. Jakarta: FKUI. 2013. p. 106-09
- [6] Havlickova B, Czaika VA, Friedrich M. Epidemiological trends in skin mycoses worldwide. *Journal compilation Blackwell Publishing Ltd.* 2008;51(4): 2-15.
- [7] Hiok-Hee T. Superficial Fungal Infections Seen at the National Skin Centre, Singapore. *Jpn J Med Mycol.* 2005;46:77-80.

- [8] Rahman MH, Hadiuzzaman Md, Jaman MK, Bhuiyan, Islam N, Ansari NP, et al. Prevalence of superficial fungal infection in the rural of Bangladesh. *Iran J Dermatol.* 2011;14: 86-91
- [9] Davis JD. Superficial fungal infection of the skin: tinea corporis, tinea pedis, and candida intertrigo. *Infectious diseases update* 1995;2:157-61.
- [10] Goodall I. HbA1c Standardisation destination-global IFCC standardization how, why, where, and when. *Clin Biochem Rev* 2005;26(1).
- [11] Ragnunatha S, Anitha B, Inamadar C, Palit A, Devarmani SS. Cutaneous disorders in 500 patients attending diabetic clinic. *Indian J Dermatol* 2011;56:2-6.
- [12] Suheyla S. The frequency of superficial fungal infection in patients with diabetes mellitus and its relation to the level of HbA1c. *Turkish J Endocrinol Metab* 1998;2:21-3.
- [13] Perkumpulan Endokrinologi Indonesia. Dalam: Konsensus pengelolaan dan pencegahan diabetes mellitus tipe 2 di Indonesia. Jakarta: PB PERKENDI; 2008.h.4-27.
- [14] Mahler RJ, Adler ML. Type 2 diabetes mellitus : Update on diagnosis, pathophysiology, and treatment. *Jcem. Endojournals.* 1999;84:1165-71.
- [15] Wani MA, Hasan I, Hayat BM, Ahmed QM. Cutaneous manifestation of diabetes mellitus: A hospital based study in Khasmir, India. *Edog* 2009;5(2).
- [16] Jahangir MU, Ahmed I, Nasreen S. Cutaneous manifestations of diabetic foot. *Jornal of Pakistan association of dermatologists* 2009;19: 13-7.
- [17] Mardila F. Hubungan Kandidiasis Intertriginosa dan Diabetes Melitus Tipe 2 di Poliklinik Kulit Dan Kelamin RSUD Dokter Soedarso. [Skripsi]. Pontianak: Universitas TanjungPura; 2012.
- [18] Badan Penelitian dan Pengembangan Kesehatan Departemen Kesehatan Republik Indonesia. Riset Kesehatan Dasar 2007. Jakarta : Departemen kesehatan RI; 2008.
- [19] Maulina, Muis K, Jusuf NK. Hubungan Kadar Hemoglobin Glikosilat (HbA1C) dengan Kandidiasis Kutis pada Pasien Diabetes Melitus Tipe 2 Di Rumah Sakit Umum Pusat Haji Adam Malik. Medan. 2014.
- [20] Wowor SR, Pandelege H.E.J, Kapantow MG. Profil Kandidosis Intertrogenosa di Poliklinik kulit dan kelamin BLU RSUP Prof. DR. D. Kandau. Manado. 2012
- [21] Armi. Pengaruh Pendidikan Kesehatan Terhadap Tingkat Kepatuhan Pasien Diabetes Melitus Dengan Retinopati Diabetik Dalam Melakukan Pemeriksaan Mata Di Rumah Sakit Aini, Jakarta. 2014.
- [22] Mongisidi, G. Hubungan Antara Status Sosio-Ekonomi Dengan Kejadian Diabetes Melitus Tipe 2 di Poliklinik Interna BLU RSUP Prof. Dr. R. D, Kandou. Manado. 2014.
- [23] Ramadona, A. Pengaruh Konseling Obat Terhadap Kepatuhan Pasien Diabetes Melitus Tipe 2 di Poliklinik Khusus Rumah Sakit Umum Pusat DR. M. Djamil Padang. Artikel. Padang: Universitas Padang. Hlm 2-3, 10-13. 2011.
- [24] Komariah, Sjam R. Kolonisasi Candida dalam Rongga Mulut. FKUI.2012.
- [25] Wahyuningsih, R, dkk. Identifikasi Candida spp. dengan Medium Kromogenik. Departement Parasitologi FK UI. 2012.
- [26] Schteingart DE. Metabolisme Glukosa dan Diabetes Melitus. Volume 1. Edisi ke-6. Di Dalam: Price SA dan Lorraine MW. Patofisiologi: Konsep Klinis Proses-Proses Penyakit. Jakarta: EGC.2006.p. 1259-70.
- [27] Shah, BR., Hux, JE. 2003. Quantifying The Risk Of Infection Disease For People With Diabetes. *Diabetes Care* 26, 510-513.