

# Serum 25-Hydroxyvitamin D Levels as a Predictive Factor for Allergic Rhinitis

Lia Restimulia<sup>\*1</sup>, Delfitri Munir<sup>1</sup>

<sup>1</sup>Department of Otorhinolaryngology Head and Neck Surgery, Faculty of Medicine, Universitas Sumatera Utara, Jl. Prof. Dr. Mansur No. 9, Medan, Indonesia

**Abstract. Introduction:** Allergy and 25-Hydroxyvitamin D deficiency were mentioned to be correlated. Still, there was limited number of studies in determining serum 25-Hydroxyvitamin D levels as a predictive factor, particularly in allergic rhinitis (AR). This study has never been implemented in Indonesia which the main objective was to explore this topic. **Methods:** Consecutive sampling was used in recruiting the fifteen subjects. Serum 25-Hydroxyvitamin D levels were quantified using electrochemiluminescence immunoassay (ECLIA) method while the total nasal symptom scores were acquired by enumerate all nasal symptoms. Data of serum vitamin D levels and TNSS were analysed statistically. **Results:** Based on our previous study, the predictive value was illustrated from linier regresion =  $10,230 - 0,281$  (vitamin D), with negative moderate correlation ( $r = -0,613$ ). This result can be comparable for someone with vitamin D level of 5.66 ng/mL that could predicted as 8.64 nasal score. **Conclusion:** 25-Hydroxyvitamin D could be a predictive factor for allergic rhinitis with negative moderate correlation. Hence, in the future research needs to be implemented.

**Keyword:** Allergic rhinitis, vitamin D, 25-hydroxyvitamin D, allergy, total nasal symptom score, predictive factor.

Received 14 September 2021 | Revised 26 October 2021 | Accepted 28 October 2021

## 1 Introduction

Allergic Rhinitis (AR) was affected 40% in total population and keeps increasing, which is caused by nasal inflammation as the result of allergen exposure [1]. Moderate to severe AR presents in 67.5% worldwide and fazed the quality of life [2].

Family history is one of the commonly used AR as predictive factors, but it is subjective and has 48-67% specificity, 22-72% sensitivity and a predictive value of less than 40%[3]. Total IgE examination has 60-70% specificity, and 26-47% sensitivity, but it cannot predict the severity of AR, while examination of the levels

---

<sup>\*</sup>Corresponding author at Faculty of Medicine, Universitas Sumatera Utara, Jl. Prof. Dr. Mansur No. 9, Medan, Indonesia.  
E-mail address: : lia\_resti@yahoo.com

of the immunological components in the placenta requires two or three types of tests to increase the predictive value[4].

Recent studies point out to deal with the immunomodulatory role of 25-hydroxyvitamin D that related to AR [5]. Epidemiological data reported a significant relationship between vitamin D and autoimmune, cardiovascular, asthma, cancer, and allergies [6]. Low vitamin D levels are act as predictive factor for atopy at 6 years child and asthma at 14 years patients [7]. Indonesia with a tropical climate exposed to the sun all year round, but the results of the study found that 25-hydroxyvitamin D deficiency rates were 35% in women aged over 65 years, 60% in young adult women, and 78.3% in children [8,9]. This high rate can be caused by many factors that affect blood 25-hydroxyvitamin D levels [8]. However, vitamin D deficiency as the prediction factor in AR in Indonesia has not been found yet [10]. Based on this explanation, the research was intended to discover serum 25-hydroxyvitamin D levels as the predictive factor for AR according to TNSS.

## 2. Methods

### *Study design*

Consecutive sampling was used in recruiting the fifteen subjects. Beforehand, the study has been approved by the Committee of Hospital Medical Research Ethics with informed consent prior to the study. The exclusion criteria such as the infection and other nasal obstruction diseases like we mentioned before.

### *Measurement*

Serum 25-Hydroxyvitamin D levels were quantified using electrochemiluminescence immunoassay (ECLIA) method. The total nasal symptom scores were acquired by enumerate all nasal symptoms. Data of serum vitamin D levels and TNSS were analysed statistically from our previous study [11].

### *Statistical Analysis*

The correlation among Serum 25-Hydroxyvitamin D levels and AR as well as the cut-off points were already obtained by Pearson correlation test and ROC curve. Linier regression was used to analyze whether vitamin D has a role as predictive factor for AR.

### 3. Results

The data of patients is tabulated in Table 1.

Table 1. Demographic Characteristic and the Relationship

Variable	Subjects (n=15)	
	Mean	SD
Age	28.87	9.01
	N	%
<b>Characteristics</b>		
Asthma	5	16.67
Dark skin	9	66.67
Body Mass index>30	2	6.67
Allergic dermatitis	6	20
Smoking	4	26.67
<b>AR Classification</b>		
Mild intermittent	1	6.67
Moderate to severe inter-mittent	1	6.67
Mild persistent	4	26.66
Moderate to severe persistent	9	60.00




Figure 1. Linier Regression.

**Table 2.** The Correlation Between Serum 25-Hydroxyvitamin D levels and TNSS.

TNSS	<i>n</i>	25-hydroxyvitamin D level		Statistical analysis	<i>p</i>
		Mean	SD		
4-6	5	11.87	7.14	<i>r</i> = -0.80*	0.000
7-9	6	8.74	3.56		
10-12	4	6.29	2.61		

\*Linier regression test

Our previous study turned up the strong negative relationship with the cut-off points related to AR was 12.83 ng/mL. Linier regression test found the prediction result was illustrated from linier regresion (Figure 1) = 10,230 – 0,281 (25-Hydroxyvitamin D), with negative moderate correlation (*r* = -0,613). From this result, 25-Hydroxyvitamin D level of 5.66 ng/mL can be analogous with predicted 8.64 nasal score.

#### 4. Discussion


This study obtained a significant divergence of serum 25-Hydroxyvitamin D levels between the AR and non-AR patients, likewise a study by Yalcinkaya et al. The average of serum 25-Hydroxyvitamin D levels in the AR group was 15.39 ng/mL while in the non-AR group was 53.80 ng/mL. Moreover, a study by Modh et al prevailed that 25-Hydroxyvitamin D deficiency turned out in 91% of the total AR samples with a significant recovery after vitamin D augmentation [10,12].

Based on the explanation above, the refinement of TNSS was happened as an immunomodulatory outcome of 25-Hydroxyvitamin D on the immune system. Milovanovic et al. stated that serum 25-Hydroxyvitamin D levels was a significant negative relationship with imunoglobulin E [13]. It was strengthen with the study implemented by Yip et al. that 25-Hydroxyvitamin D could repress the mast cells activity [14]. Furthermore, Vasiliou et al. found related result that serum 25-Hydroxyvitamin D levels were have relationship with asthma [15]. In addition, the escalate mice serum 25-Hydroxyvitamin D levels after UV-B exposure was related with reduction of airway inflammation compared with the non exposed to UV-B mice [16,17].

There are many factors that influence 25-Hydroxyvitamin D level, such as genetic factor , life style that tend to be indoor activities and smoking, climate/atmosphere, age, body mass index, skin pigmentation, and also the time and

duration of sun exposure [18]. Allowing that the previous archetype claimed that 25-Hydroxyvitamin D was completely metabolized from the sun and food intake in kidney, discovery now stated that the production of 25-Hydroxyvitamin D could be generated by some types of immune cells with vitamin D receptor (VDR), such as dendritic cells and T-cells. These cells along with B cells and mast cells could expressed this VDR. The serum 25-Hydroxyvitamin D produced by genetic that regulates two thousand different genes, which one of them is a gene that took parts in allergies [19]. This vitamin encodes the related to allergic rhinitis and asthma chromosome, namely chromosome 12. This vitamin regulates chromosomes influenced by IgE, 13Q14 and 7Q14. This hypothesis explains how genetic factors have the the main role for the serum 25-Hydroxyvitamin D levels in AR patient [16,18].

This study indicated a strong negative correlation between serum 25-Hydroxyvitamin D levels with AR incidence and TNSS, which showed that 25-Hydroxyvitamin D plays an important role in the AR. Moreover, this study were in accordance with Thakkar et al. with negative correlation between serum 25-Hydroxyvitamin D levels with TNSS with a moderate relation strength [20,21].



**Figure 2. Vitamin D regulation in Allergic rhinitis.** Vitamin D regulate the proinflammatory cells by inducing IL-10 and T regulator cell as well suppressing pro inflammatory cytokines.

Vitamin D has the capability in controlling regulation of inflammatory Th2-mediated cell, namely the dendritic cell by reducing differentiation and

lipopolysaccharide activity (LPS) [22]. Furthermore, it can reduce differentiation of mast cells and induce its apoptosis in 30-40 days by suppressing Immunoglobulin-E and IL-4 [15]. A research by Yip et al. found the inhibition of mast cells degranulation in mice after 25-Hydroxyvitamin D supplementation [14]. The study by Hypponen et al. stated that serum 25-Hydroxyvitamin D levels had a significantly negative relationship with Immunoglobulin-E [23]. Histamine-receptor binding inhibition in the nasal mucosa causes the suppression of Th2 activity in the early stage. Induction of interleukin 10 as anti-inflammatory is influenced by APC, Th2 cells, and mast cells and can also play a role in Th2 suppression (Figure 2) [23].

Still, in restraining the regulation of Th-2 inflammatory cell, 25-Hydroxyvitamin D can impede the differentiation and proliferation of B-cells become plasma cells by increasing the work of IL-10 and suppressing the pro-inflammatory cytokines and chemokines action [22]. The 25-Hydroxyvitamin D can reduce the recruitment of blood eosinophils into mucosa in late phase, consequently decreasing the differentiation of B-cells into plasma cells. Inhibition of IL-5 by 25-Hydroxyvitamin D delays the process of eosinophilic maturation, differentiation, migration, and infiltration towards nasal mucosa [14].

Vitamin D also has a role in suppressing the inflammatory responses by increasing the regulation of Th1-mediated cell, for example TGF- $\beta$ , NK cells and IL-10 [14]. It may also repress TLR and MHC class II activation by inhibiting Th1-mediated cells, such as IFN- $\gamma$ . This action inhibits proinflammatory cytokines regulation [23].

In conclusion, this result found that 25-Hydroxyvitamin D can be a predictive factor for AR. However, future randomized controlled trials are needed.

### **Acknowledgement**

The authors would like to be grateful for many supports from our lecturers

### **Funding and Conflict of Interest**

The authors did not receive funding from any institution, and there was no conflict of interest which should be reported.

## References

1. Small. P, Kim. H, “Allergic Rhinitis”. *Allergy, Asthma & Clin Immunol*, vol.14 no. 2 pp. 51. 2018.
2. Bauchau. V, Durham. S, “Prevalence and Rate of Diagnosis of Allergic Rhinitis in Europe”, *Eur Respir J*, vol. 24, no. 5, pp. 758-64. 2004.
3. Hodder. M, Tulic. M, Prescott. S, “Algorithmic models to predict allergic disease using multiple neonatal markers”, *Indian Pediatrics*, no. 46, pp291-2. 2009.
4. Nickolls. C, Campbell. D, “Top 10 food allergy myths”, *Journ Ped Child Health*, no. 1, pp. 1-5. 2015.
5. Seidman. M, Gurgel. R, Lin. S, Schwartz. S, Barody. F, “Clinical Practice Guideline: Allergic Rhinitis”. *Otolaryngol Head and Neck Surg*, vol. 152, no. 1, pp 1-432015. 2015.
6. Neubauer. P, Lauster. R, Worm. M, Garbe. L, “Vitamin D receptor activation modulates the allergic immune response”, *University of medicine, Berlin*. 2011.
7. Hollams. E, Hart. P, Holt. B, Serralha. M, Parson. F, Klerk. N, *et al.*, “Vitamins D and atopy and asthma phenotypes in children: a longitudinal cohort stud”, *Eur Respir J*, no.38, pp. 1320-7. 2011.
8. Arabi. A, Rassi. R, Fuleihan. G, “Hypovitaminosis D in developing countries- prevalence, risk factors and outcomes”, *Endocrinol*, no. 6, pp.550-612010.
9. Soesanti. F, Pulungan. A, Tridjaja. B, Batubara. J, “Vitamin D profile in healthy children aged 7-12 years old in Indonesia”, *Int J Pediatr Endocrinol*, pp 167. 2013.
10. Modh. D, Katarkar. A, Thakkar. B, Jain. A, Shah. P, Joshi. K, “Role of Vitamin D Supplementation in Allergic Rhinitis”. *Indian Journal of Allergy*, vol. 28, no. 1, pp 35-9. 2014.
11. Bousquet. J, Khaltaev. N, Cruz. A, Denburg. J, Fokkens. W, Togias. A, *et al.*, “Allergic Rinitis and its Impact on Asthma (ARIA) 2008 Update (in Collaboration with the World Health Organization)”. *Journal Allergy*. Vol. 63, no. 86, pp. 8–160. 2008.
12. Yalcinkaya. E, Tunckasik. M, Guler. I, Kocaturk. S, Gunduz. O, “Evaluation of The Correlation of 25-hydroxyvitamin-D Serum Levels with Allergic Rhinitis”. *ENT Updates*, vol. 5, no. 1, pp. 19-22. 2015.
13. Milovanovic. M, Heine. G, Hallatschek. W, Opitz. B, Radburch. A, Worm. M, “Vitamin D receptor binds to the  $\epsilon$  germline gene promotor and exhibits transrepressive activity”, *J Allergy Clin Immunol*, vol. 126, no. 5, pp. 1016-23. 2010.

14. Yip. K, Kolesnikoff. N, Yu. C, Hauschild. N, Taing. H, “Mechanisms of Vitamin D3 Metabolite Repression of IgE-Dependent Mast Cell Activation”. *J Allergy Clin Immunol*, vol. 133, no. 5, pp. 1356-64. 2014.
15. Vasiliou. J, Lui. S, Chohan. V, Xystrakis. E, Bush. A, Hawrylowicz. C, “Vitamin D Deficiency Induces Th2 Skewing and Eosinophilia in Neonatal Allergic Airways Disease”, *Allergy*, vol. 69, no. 5, pp. 1380-9. 2014.
16. Lange. N, Litonjua. A, Hawrylowicz. C, Weiss. S, “Vitamin D, the Immune System and Asthma”, *Expert Rev Clin Immunol*, vol. 95, no. 6, 693-702. 2009.
17. Hulley. S, Cummings. S, Browner. W, Grady. D, Newman. T, “Designing Cross-Sectional and Case-Control Studies”. In: Hulley S, eds. *Designing Clinical Research*, 3<sup>th</sup> edition, Philadelphia: Lippincott Williams & Wilkins, pp. 109-25, 2007.
18. Lucock. M, Jones. P, Martin. C, Hons. B, Beckett. E, Yates. Z, et al., “Vitamin D: Beyond Metabolism”, *Evidence-Based Complementary & Alternative Medicine*, vol. 20, no. 4, pp. 310-22. 2015.
19. Cheng. H, Kim. S, Park. G, Chang. S, Bang. S, Won. C, et al., “Low Vitamin D Levels are Associated with Atopic Dermatitis, but Not Allergic Rhinitis, Asthma, or IgE Sensitization, in the Adult Korean Population”. *J Allergy Clin Immunol*, vol. 133, no. 4, pp. 1048-55. 2014.
20. Restimulia. L, Pawarti. D, Mayaguyang. H., “The Relationship between Serum Vitamin D Levels with Allergic Rhinitis Incidence and Total Nasal Symptom Score in Allergic Rhinitis Patients”, *Open Acces Med J Med Sci*, vol. 6, no. 8, pp. 1405-09. 2018.
21. Thakkar. B, Katarkar. A, Modh. D, Jain. A, Shah. P, Joshi. K, “Deficiency of Vitamin D in Allergic-Rhinitis: a Possible Factor in Multifactorial Disease”, *J Clin Rhinol*, vol. 7, no. 3, pp. 112-6. 2014.
22. Mulligan. J, White. D, Wang. E, Sansoni. S, Moses. H, Yawn. R., et al., “Vitamin D3 Deficiency Increases Sinus Mucosa Dendritic Cells in Pediatric Chronic Rhinosinusitis with Nasal Polyps”, *Otolaryngol Head Neck Surg*, vol. 147, no. 4, pp. 773-81. 2012.
23. Hypponen. E, Berry. D, Wjst. M, Power. C, “Serum 25-Hydroxyvitamin D and IgE- a Significant but Nonlinier relationship”, *Allergy*, no. 64, pp. 613-20. 2009.