

Relationship between Allergic Rhinitis to Quality of Life of Students of Faculty of Medicine Universitas Sumatera Utara Batch 2018

Anna Delinda^{1*}, *Andrina Rambe*²

¹Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia

²Department of Ear Nose and Throat, Universitas Sumatera Utara, Medan, Indonesia

Abstract. Background: Allergic rhinitis is an inflammation mediated by Immunoglobulin E (IgE) with symptoms such as runny nose, sneezing, nasal congestion, with or without itchy nose. Although allergic rhinitis is not a dangerous disease, this disease has an impact on decreasing the quality of life of the sufferer which can interfere with daily activities. **Objective:** This study use to determine the relationship between allergic rhinitis and quality of life among students of the Faculty of Medicine, Universitas Sumatera Utara, batch 2018. **Methods:** This study used an observational analytical method with a cross-sectional study approach. The research sample was determined by simple random sampling technique and used two questionnaires, namely SFAR to assess allergic rhinitis and SF 36 to assess quality of life. **Results:** Based on the study, the number of allergic rhinitis sufferers from 72 students of the Faculty of Medicine, Universitas Sumatera Utara, batch 2018, 38 students suffering from allergic rhinitis. The results of data analysis using the Chi Square statistical test showed that there was a significant relationship between allergic rhinitis and quality of life, p (p value) of 0.001 ($p < 0.05$). **Conclusion:** There is a relationship between allergic rhinitis and quality of life in 2018 students of the Faculty of Medicine, Universitas Sumatera Utara.

Keyword: Allergic rhinitis, Quality of life, SFAR, The Short Form-36

Received 02 January 2022 | Revised 02 March 2023 | Accepted 02 March 2023

Introduction

According to *World Health Organization Allergic Rhinitis and its Impact on Asthma* (WHO ARIA), allergic rhinitis is an inflammation of the nasal mucosa after exposed to IgE-mediated allergens, showed by nasal symptoms, such as, *rhinorrhea* (watery runny nose), sneezing, nasal congestion and itchy nose [1].

The prevalence of allergic rhinitis is increasing progressively in developed country, especially adults, 10%-40%, on the other side, rate for the children is 2%-25% worldwide. Economic

*Corresponding author at: Universitas Sumatera Utara, Medan, Indonesia

E-mail address: [author's email address]

development, westernized lifestyle, urbanization and eating practice also increase the case of allergic rhinitis in developing countries [2].

It is acknowledged that allergic rhinitis have an impact to those sufferers. Although allergic rhinitis is not a dangerous disease, but this can also affect on decreasing the quality of sufferers' life – their daily life [3].

In a study involving 1.948 individuals who completed RQLQ questionnaire, there were 3 parameters which indicate the effect of allergic rhinitis, such as embarrassment, frustration, and practical problems, also include discomfort to carrying tissue or handkerchief, the need to wipe the nose, produce nasal snot continuously [4].

There are several instruments that can be used to assess the quality of life of patients with allergic rhinitis. SF-36 is an instrument which widely used by researchers. This instrument is quick (5-10 minutes) and easy to use, also can be done using a telephone interview [5].

Therefore, based on the background displayed above, the writer is interested in knowing the relationship between allergic rhinitis and the quality of life of students of the Faculty of Medicine, Universitas Sumatera Utara, batch 2018.

Methodology

The research design used in this study was *observational analytic* study with *cross-sectional* study approach. This study intend to determine the relation between allergic rhinitis to the quality of life of student of the Faculty of Medicine, Universitas Sumatera Utara, batch 2018. This study was conducted by communication network of Faculty of Medicine, Universitas Sumatera Utara, throughout July to November in 2021. The population in this study were all students of the Faculty of Medicine, Universitas Sumatera Utara, batch 2018 – numbered 254 students.

The number of samples will be calculated using the formula:

$n = 1 + Ne_2$ Description :

n = sample N

= population e

= error tolerance (0,1)

Based on this formula, the minimum number of sample is:

$$n = 1 + \frac{254}{(0,1)^2}$$

$$n = \frac{254}{3,54} = 72$$

This study needed 72 students as sample. *Simple random sampling* technique used to determine randomly the student of Faculty of Medicine, Universitas Sumatera Utara which process will give every student (population) and equal chance of being sampled.

Result

From the collection of respondents using an online questionnaire, the authors collected 72 respondents who participated in this study. The subjects of this study were students of the Faculty of Medicine, Universitas Sumatera Utara, batch 2018. From the data, the distribution of respondents was obtained based on the diagnosis of allergic rhinitis.

Table 1. Frequency distribution of respondents based on the diagnosis of allergic rhinitis

Diagnosis of Allergic Rhinitis	n	%
Students Not Suffering From Allergic Rhinitis	34	47,2%
Students with Allergic Rhinitis	38	52,8%
Total	72	72 (100%)

Table 1 shows that the respondents were divided into two groups, there were 38 students with allergic rhinitis (52.8%) and 34 students weren't rhinitis sufferers (47.2%).

Table 2 Distribution of allergic rhinitis by sex

Sex	Students Are Not Suffering From Allergic Rhinitis		Students with Allergic Rhinitis		
	n	%	n	%	
Male	10	35,7%	18	64,3%	28
Female	24	54,5%	20	45,5%	44
Total	34		38		72

From table 2, it can be seen that the distribution of respondents by gender shows that the amount of female allergic rhinitis sufferers is more than male sufferers. For female the amount is 20 people (45.5%), while the male is 18 (64.3%). For not sufferers, the percentage for female is 24 people (54.5%) and male is 10 people (35.7%).

Table 3 Distribution of allergic rhinitis by family history

Family History	Students Are Not Suffering From Allergic Rhinitis	Students with Allergic Rhinitis	Students (n) %
Without family history	29	14	43 (59,7%)
With family history	5	24	29 (40,3%)
Total	34	38	72 (100%)

The data from table 3 shows that the distribution by family history of disease was 72 respondents with 29 respondents of students who have family history of allergic rhinitis (40.3%). From that category, there are 5 students are not sufferers and 24 students are the sufferers. The other category, there are 43 respondents (59.7%) without a family history of disease with 14 people with allergic rhinitis and 29 non-allergic rhinitis sufferers.

Table 4 Distribution of allergic rhinitis based on clinical symptoms of rhinitis

Rhinitis Clinical Symptoms	Students Suffering From Allergic Rhinitis (n)	Not Suffering with Allergic Rhinitis (n)	Students (n) %
Asymptomatic	6	5	11 (15,3%)
sneeze	8	5	13 (18,1%)
Runny nose	0	1	1 (1,4%)
Nasal congestion	4	0	4 (5,5%)

Itchy nose & watery eyes	1	0	1 (1,4%)
Sneezing & runny nose	3	0	3 (4,2%)
Sneezing & nasal congestion	2	1	3 (4,2%)
Sneezing, itchy nose & watery eyes	0	3	3 (4,2%)
Runny nose & nasal congestion	0	1	1 (1,4%)
Nasal congestion, itchy nose & watery eyes	0	3	3 (4,2%)
Sneezing, runny nose & nasal congestion	5	5	10 (13,8%)
Sneezing, runny nose, itchy nose & watery eyes	0	1	1 (1,4%)
Sneezing, nasal congestion, itchy nose & watery eyes	0	1	1 (1,4%)
Sneezing, runny nose, nasal congestion,	5	12	17 (23,5%)

itchy nose & watery eyes			
Total	34	38	72 (100%)

Table 4 shows that the most frequent clinical symptoms experienced by respondents are sneezing, runny nose, nasal congestion, itchy nose & watery eyes with total 17 respondents (23.5%), consisting 12 people with allergic rhinitis and 5 people who are not allergic rhinitis sufferers. This was followed by the symptoms of sneezing with 13 respondents (18.1%) consisting of 5 people with allergic rhinitis and 8 people not suffering from allergic rhinitis.

Table 5 Distribution of allergic rhinitis by history of other atopic diseases

History of Atopic Disease	Students Suffering From Allergic Rhinitis (n)	Not Allergic Rhinitis (n)	Students with Allergic Rhinitis (n) %
No comorbid	22	18	40 (55,5%)
Asthma	5	6	11 (15,3%)
Eczema	1	1	2 (2,8%)
Allergic rhinitis	5	8	13 (18,1%)
Asthma & allergic rhinitis	1	3	4 (5,5%)
Eczema & allergic rhinitis	0	1	1 (1,4%)
Asthma, eczema & allergic rhinitis	0	1	1 (1,4%)
Total	34	38	72 (100%)

Table 5 represents data of distribution of family history of other atopic diseases with allergic rhinitis is the most disease experienced by the respondents with total 13 people (18,1%) consisting 8 people with allergic rhinitis and 5 people are not the sufferers. The other disease is asthma with total 11 people (15,3%), consists 6 people with allergic

rhinitis and 5 people are not the sufferers. From the family with asthma and allergic rhinitis total 4 people (5,5%), contain 1 people not the sufferer and 3 people with allergic rhinitis.

Table 6 Distribution of respondents by quality of life

Quality of life	n	%
Poor	20	27,8%
Good	52	72,2%
Total	72	100%

Table 6 shows that the respondents with good quality of life are 52 people (72.2%) and 20 people (27.8%) have poor quality of life.

Table 7 Distribution of allergic rhinitis respondents based on quality of life

Quality of life	Students	Students	Students (n)
	Not Suffering From Allergic Rhinitis (n)	with Allergic Rhinitis (n)	%
Poor	3	17	20 (27,8%)
Good	31	21	52 (72,2%)
Total	34	38	72 (100%)

From table 7 the distribution of respondents with good quality of life shows that 52 people (72.2%) with 21 are sufferers of allergic rhinitis and 31 people are not the sufferers from allergic rhinitis. While respondents with poor quality of life were 20 people (27.8%) with 17 people suffering from allergic rhinitis and 3 people not suffering from allergic rhinitis.

Table 8 Cross-tabulation of allergic rhinitis with quality of life

		Quality of life				P value
		Poor		Good		
Allergic rhinitis	No	3	15%	31	59,6	0.001
	Yes	17	85%	21	40,4%	
Total		20	100	52	100	

From table 8 it can be seen that the p value is 0.001 ($p < 0.005$), which indicates that there is a relationship between allergic rhinitis and the quality of life of the sufferer.

Dicussion

The purpose of this study is to determine the relationship between allergic rhinitis and quality of life of sufferers. Based on this study, the p value was 0.001 ($p < 0.005$). From the results, can be concluded that there is a relationship between allergic rhinitis and the patient's quality of life in students of the Faculty of Medicine, Universitas Sumatera Utara.

The results of this study are in line with research by Novian (2016) where the p value - 0.022 ($p < 0.05$) states that there is a significant difference in the quality of life between the positive allergic rhinitis group and the negative allergic rhinitis group [10]. In general, patients are bothered by the symptoms that occur on the nose (nose obstruction, rhinorrhea and sneezing). They experience discomfort carrying tissue or handkerchief, the need to wipe the nose or eyes, and persistent nasal discharge. They are also disturbed by not being able to sleep well at night and often become tired, irritable, memory impaired, and sleepy during the day. They experience limitations in daily activities, which makes them frustrated and distracted [4].

Penelitian ini dilakukan uji hipotesis dengan metode Chi Square dengan tingkat kemaknaan 0,05 ($\alpha=5\%$) diperoleh nilai p (p value) sebesar 0,001 ($p<0,05$) yang berarti hasilnya signifikan. Oleh karena itu, hipotesis H_0 ditolak. Hal tersebut bahwa ada hubungan antara penyakit rinitis alergi dengan kualitas hidup pada mahasiswa Fakultas Kedokteran Universitas Sumatera Utara.

This research was conducted to test the hypothesis using the Chi Square method with a significance level of 0.05 ($\alpha = 5\%$) and the p value was 0.001 ($p < 0.05$), which indicates significance. Therefore, the hypothesis H_0 is rejected. This means that there is a relationship between allergic rhinitis and quality of life in students of the Faculty of Medicine, Universitas Sumatera Utara.

Conclusion

There is a significant relationship between allergic rhinitis and the quality of life of students of the Faculty of Medicine, Univeristas Sumatera Utara batch 2018 statistically where the p-value is 0.001 ($p < 0.05$).

Reference

- [1] Brożek, J.L., Bousquet, J., Agache, I., Agarwal, A., Bachert, C., et al. 2017, 'Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines—2016 revision', *Journal of Allergy and Clinical Immunology*, 140(4), pp.950-958.
- [2] Zhang, L., & Zhang, Y., 2019. Increasing prevalence of allergic rhinitis in China. *Allergy, Asthma and Immunology Research*, 11(2), 156–169. <https://doi.org/10.4168/aaair.2019.11.2.156>
- [3] Meltzer EO, Diego S. 2011. Quality of life in adults and children with allergic rhinitis. *J. Allergy Clin. Immunol.* 108(1):45-53.
- [4] Cristina Camelo-Nunes, I. and Solé, D. 2010 *Allergic rhinitis: indicators of quality of life* Rinite alérgica: indicadores de qualidade de vida, J Bras Pneumol.*
- [5] Tinartayu, S., & Riyanto, B. U. D. (2015). SF-36 sebagai instrumen penilai kualitas hidup penderita tuberkulosis (TB) paru. *Mutiara Medika: Jurnal Kedokteran Dan Kesehatan*, 15(1), 7–14. <https://journal.umy.ac.id/index.php/mm/article/view/2488/2552>
- [6] David, Sigarlaki. 2017. Perbedaan Kualitas Hidup Penderita dan Bukan Penderita Rinitis Alergi Pada Mahasiswa Fakultas Kedokteran Universitas Lampung Tahun 2016.
- [7] Nadira, Alya. 2020. Hubungan Rinitis Alergi dengan Depresi Pada Siswa Pondok Pesantren Darul Ulum Banda Aceh.
- [8] Yuksel H, Sakar A, Yilmaz O, Yorgancioglu A, Celik P, Ozcan C. 2008. Prevalence and comorbidity of allergic eczema, rhinitis , and asthma in a city in western turkey. *J Investig Allergol Clin Immunol.* 18(1):31-5.
- [9] Nurjannah. 2011. Faktor risiko rinitis alergi pada pasien rawat jalan di poliklinik THT- KL rumah sakit umum daerah zainoel abidin (RSUDZA) banda aceh tahun 2011. *Jurnal Kedokteran Syiah Kuala.* 11(2):60-5.
- [10] Novian, Maya. 2016. Hubungan Rinitis Alergi Dengan Kualitas Hidup Mahasiswa Fakultas Kedokteran Universitas Sumatera Utara.